



Dynamic Interventions: Opportunities and Challenges

Bonnie Spring
Center for Behavior and Health
Northwestern University



Optimizing Obesity Treatment: 1 Static and 2 Dynamic Examples

- 1. MOST:** Multiphase Optimization Strategy
- 2. SMART:** Sequential Multiple Assignment Randomized Trial
- 3. JITAI:** Just in Time Adaptive Intervention

Opt-In Study

Optimization of Remotely Delivered Intensive Lifestyle Treatment for Obesity

Principal Investigators

Bonnie Spring, Ph.D. (Northwestern University)

Linda Collins, Ph.D (Pennsylvania State University)



NIDDK R01 DK097364



Opt-IN MOST Design – Optimize Components for Best Average Fixed Treatment

- ▶ Decision point – outset (choose components for fixed intervention)
- ▶ Intervention options – components
- ▶ Tailoring variable
 - ▶ Weight loss
 - ▶ Cost
 - ▶ [Reach]
- ▶ Decision rule – optimize for specific constraints
 - ▶ Max weight loss for \$500
 - ▶ [Max enrollment]

Component	Rank Cost	Rank Engagement
Texting	1	2
PCP updates	2	5
Buddy training	3	3
Coaching	4	4
Meal replacement	5	1

SMART Weight Loss Management – Optimize Best Treatment Sequence and Tactic for Addressing Nonresponse

- ▶ **Decision Points:**
 - ▶ Outset - Best first line treatment
 - ▶ 2 weeks, 1 month – Optimal tactic to address treatment nonresponse
- ▶ **Intervention Options:**
 - ▶ Components: Text, Coaching, Meal replacement
- ▶ **Tailoring variable**
 - ▶ Weight loss
- ▶ **Decision rule – Adaptive:** do one thing if responds, another if not
- ▶ **Once optimized, decision rule/algorithm remains the same**
 - ▶ If response, continue
 - ▶ If nonresponse – continue (More) or Augment

JITAI – Optimize intervention to particular person and their changing needs over time

- ▶ Decision points – continuous patient data; intervene any time
- ▶ Intervention options –
 - ▶ Digital, mobile – text, coach call, call or be called by buddy – (MR “piggybank”)
- ▶ Tailoring variables – Many continuously available
 - ▶ Objective sensor data on energy absorption, physical activity, sleep, stress, social activity
- ▶ Decision rules – Dynamic – push, pull, learn not fixed and only push out. Pull in and do machine learning about person’s response over time (follow text suggestion? Take call from coach or friend?)
 - ▶ Can take contextual information into account – cues, access, others in room




Dynamic Intervention Opportunities

- ▶ Continuous data – learn more comprehensively and faster;
 - ▶ develop better treatment algorithms
 - ▶ Different information channels – EMA, physiological, environmental cues, location
- ▶ Intervene in real time, when needed, just in time
- ▶ Personalized treatment algorithms!
- ▶ Understand/develop theory of how mediating pathways change over time

Dynamic Intervention Challenges

- ▶ Too much information!!
 - ▶ How to capture, visualize, and make the data deluge actionable?
- ▶ Lack of dynamic behavioral science theory that goes beyond snapshots
- ▶ Team science interdisciplinary challenges: what's a measure?
- ▶ Systems integration with EHR, health care work flow



Thank you!

bspring@northwestern.edu