

Appendix B

Global Smokeless Tobacco Product Factsheets

Chapter Contents

African Traditional Snuff Products	B-3
Betel Quid With Tobacco	B-5
Chimó	B-7
Creamy Snuff	B-9
Dissolvable Tobacco	B-11
Dohra	B-13
Dry Snuff	B-15
Gudakhu	B-17
Gul	B-19
Gutka (Pan Masala with Tobacco)	B-21
Iqmik	B-23
Khaini	B-25
Kiwam	B-27
Loose Leaf Chew	B-29
Mainpuri	B-31
Mawa	B-33
Mishri	B-35
Moist Snuff (Dip)	B-37
Nass	B-39
Plug	B-41
Rapé	B-43
Red Toothpowder	B-45
Shammah	B-47
Snus	B-49
Tapkeer	B-51
Tobacco Leaf	B-53
Tobacco Water (Tuibur)	B-55
Tombol	B-57
Toombak	B-59
Twist	B-61
Zarda	B-63

African Traditional Snuff Products

A wide range of smokeless tobacco products are used in the 46 countries that make up the WHO African Region. Smokeless tobacco products can take a variety of forms, are available commercially, and can be custom-made or manufactured in small factories. The products can be chewed, sniffed, sucked, or applied to teeth and gums. Several of the products are summarized here.



Photo courtesy of Olalekan Ayo-Yusuf, University of Pretoria

Product Types, Modes of Absorption, and Main Geographic Locations

Ghana traditional snuff (tawa)

This local dry snuff, often called “tawa,” is prepared by mixing the dried tobacco leaf with some chemicals such as saltpeter (potassium nitrate) and grinding it into a fine powder. It may be held in the mouth or used nasally to induce sneezing to “lighten” the head; it may also be used as a depressant or stimulant.¹

Neffa (naffa, tenfeha, nufha)

These are the names given to dry snuff that is used nasally in northern Africa. Neffa/naffa is used in Tunisia, Libya, and Algeria. Tenfeha is used in Morocco, and nufha in Algeria.²

Snuif

Snuif, the South African word for “snuff,” is not a specific type of smokeless tobacco product. Snuif is both commercially and locally produced. Some brand names are Singleton (dry snuff), Taxi, and Ntsu (moist snuff). The custom-made traditional mix is prepared by hand-mixing finely ground sun-dried tobacco leaf with ash from local plants.³

Taaba

This type of smokeless product is used in a number of West African countries, including northern Nigeria, Cameroon, Senegal, Uganda, and Chad. It is locally produced from dry fermented tobacco that is pulverized and mixed with natron/atron (a

naturally occurring mixture of sodium bicarbonate and sodium chloride). Taaba can be used nasally or sucked orally.

Prevalence and Demographics

Ghana traditional snuff (tawa)

Although no data are available specifically on the prevalence of use of traditional snuff, findings from the Global Youth Tobacco Survey (GYTS) in 2006 showed that 10.4% of youths in Ghana reported using “tobacco products other than cigarettes,” including snuff.⁴

Neffa

Neffa is predominantly used by men. Although no data are available specifically on the prevalence of neffa use, in the Algerian provinces of Oran, Constantine, and Setif, youth prevalence of tobacco use other than cigarettes was 7.8–8.9%, according to the 2007 GYTS.⁴

Snuif

Snuif is the name given to snuff in South Africa. Its use is most common among women and people living in rural areas, as well as individuals who are older, black, or have less education and income.⁵ A 2003 study in South Africa reported that 2.4% of men and 10.9% of women aged 15 years and older reported ever using any smokeless tobacco, including snuff.⁶ The 2008 South African National Youth Risk Behavior Survey found that 12.1% of adolescents used smokeless tobacco in the past month.⁷

African Traditional Snuff Products

Taaba

No data are available specifically on the prevalence of dry snuff use, but the 2008 Nigeria Demographic and Health Survey found that 3.2% of adult men and 0.5% of adult women report using smokeless tobacco.⁸ Although national prevalence data in Nigeria suggest relatively low use rates, 2007 data from a state in Nigeria's North-East geopolitical zone revealed smokeless tobacco use rates as high as

10.8% among men and 4.1% among women aged 15 years and older.⁹ The 2008 GYTS in Nigeria found that youth prevalence of tobacco use other than cigarettes was between 13.1% and 23.3% in five states.⁴ In Uganda in 2006–2007, the prevalence of any smokeless tobacco use among adults (age 15–54) was 3.9% for men and 2.6% for women,⁸ and among adolescents (age 13–15), 8.6% for boys and 9.6% for girls.⁴

Chemical Measurements

Only limited data are available on the toxicity of smokeless tobacco products used in the region, but recent data suggest considerable variability in the toxicity and nicotine profiles of some of the products that have been tested. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine mg/g wet wt	Free Nicotine mg/g wet wt	NNK	NNN	NNAL	Total TSNA ^s *
		ng/g wet wt					
Snuif, traditional snuff, South Africa	9.29	5.29	5.01	1,610	5,570	71.8	20,500
Traditional snuff, Nigeria	9.42	2.49	2.39	285	711	29.5	1,520

*Total TSNA^s represent the sum of NNK, NNN, and NNAL (shown), and *N*'-nitrosoanatabine and *N*'-nitrosoanabasine (not shown).
Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N*'-nitrosonornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNA^s = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.
Source: Stanfill et al. 2011 (10).

References

1. Addo MA, Gbadago JK, Affum HA, Adam T, Ahmed K, Okley G. Mineral profile of Ghanaian dried tobacco leaves and local snuff: a comparative study. *J Radioanal Nucl Chem.* 2008; 277(3):517–24.
2. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
3. Ayo-Yusuf OA, Peltzer K, Mufamadi J. Traditional healers' perceptions of smokeless tobacco use and health in the Limpopo Province of South Africa. *Subst Use Misuse.* 2006;41:211–22.
4. Centers for Disease Control and Prevention. [Unpublished data from the Global Youth Tobacco Survey (GYTS).] Atlanta: Centers for Disease Control and Prevention; 2008–2010.
5. World Health Organization. WHO report on the global tobacco epidemic, 2011. Appendix VIII—Table 8.2: Crude smokeless tobacco prevalence in WHO member states. Geneva: World Health Organization; 2011. Available from: http://www.who.int/tobacco/global_report/2011/en_tf_global_report_2011_appendix_viii_table_2.pdf
6. Peer N, Bradshaw D, Laubscher R, Steyn K. Trends in adult tobacco use from two South African Demographic and Health Surveys conducted in 1998 and 2003. *S Afr Med J.* 2009 Oct;99(10):744–9.
7. Reddy SP, James S, Sewpaul R, Koopman F, Funani NI, Sifunda S, et al. Umthente Uhlaba Usamila – The South African Youth Risk Behaviour Survey 2008. Cape Town: South African Medical Research Council; 2010.
8. Kishor S, et al. Prevalence of current cigarette smoking and tobacco use among women and men in developing countries. Forthcoming, 2014.
9. Desalu OO, Iseh KR, Olokoba AB, Salawu FK, Danburam A. Smokeless tobacco use in adult Nigerian population. *Niger J Clin Pract.* 2010;13(4):382–7.
10. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. *Tob Control.* 2011 May;20(3):e2. Epub 2010 Nov 25.

For additional information on African traditional snuff products, please refer to Chapter 12: Smokeless Tobacco Use in the African Region.

Betel Quid

With Tobacco

Betel quid is commonly used in many countries in the Asia-Pacific region. It can be prepared in a variety of ways depending on the region, but usually contains areca nuts, slaked lime, and catechu (extract of the *Acacia catechu* tree) wrapped in a betel leaf. Betel quid itself is not a tobacco product, but tobacco is often added to it. Chewing betel quid without tobacco is an ancient practice in India; this product is known as "tambula" in Sanskrit.¹



Photo courtesy of World Health Organization South-East Asia Regional Office and Dharendra N. Sinha

Common Names

Paan or pan (India), khilli paan (Bangladesh)

Brand Names

None

Main Geographic Locations

(WHO Region: Country)

South-East Asia Region: India, Sri Lanka, Bangladesh, Myanmar, Thailand, Indonesia, Nepal, Maldives²;

Eastern Mediterranean Region: Pakistan, United Arab Emirates^{2,3}; **Western Pacific Region:** Lao Democratic People's Republic, Palau, Cambodia, Malaysia, Vietnam, Federal States of Micronesia^{2,4}

Prevalence and Demographics

Prevalence and demographic profiles of betel quid chewers vary greatly by region. The Global Adult Tobacco Survey in India (2009–2010) shows that about 6.2% percent of all adults aged 15 years and older (7.5% of males and 4.9% of females) report using betel quid with tobacco in India,⁵ and 24.3% of all Bangladeshis aged 15 years and older (23.5% of males and 25.2% of females) consume betel quid with tobacco.⁶ The 2009–2010 Asian Betel–Quid Consortium study found that use of betel quid with tobacco among adults aged 15 and older ranges widely across several other South-East Asian countries: Prevalence is high in Nepal (males, 43.6%; females, 34.9%) and among women in Indonesia (31.7% among females, compared to 10.4% among

males). Prevalence is moderate in Malaysia (males, 6.2%; females, 12.0%), and generally low in Sri Lanka (males, 6.4%; females, 3.2%).⁷ In Myanmar in 2004, approximately 16.2% of adults aged 15 years and older (males, 27.8%; females, 4.4%) chewed betel quid with tobacco.⁸

Mode of Absorption

Oral (chewed, held in mouth)

Use Patterns

Different regions will use different types of tobacco in betel quid, such as sada pata (plain tobacco flakes) and zarda (flavored tobacco flakes) in Bangladesh and India,^{1,6} kiwam (tobacco paste) in Pakistan and India,^{1,2} and even half a cigarette in Palau.^{4,9} In Indonesia, Vietnam, and Cambodia, users may use tobacco to clean their teeth after chewing betel quid rather than inserting it directly in the quid.^{9,10} Some users swallow the juices produced from chewing betel quid.²

Main Ingredients

Tobacco, areca nut, slaked lime (calcium hydroxide) or other alkaline agents, betel leaf, and usually catechu (*Acacia catechu* tree extract). Additional ingredients vary regionally according to local preference, and can include cardamom, saffron, cloves, camphor, aniseed, turmeric, mustard, or sweeteners.^{2,9}

Betel Quid

With Tobacco

Processing/Manufacturing

Cottage industry and custom-made: Betel quid is prepared by individual vendors for sale or assembled at home by individual users. Betel quid may be prepared in a variety of ways. Areca nut can be raw, boiled, roasted, fermented, or sun-dried. Tobacco may be used raw, sun-dried, or roasted, and then finely chopped or powdered. Alternatively, the tobacco may be boiled with

molasses and made into a paste. The tobacco may then be perfumed or flavored. Slaked lime and sometimes catechu are smeared on a betel leaf, then the betel leaf is folded into a funnel shape, and tobacco, areca nut, and any other ingredients are added. The top of the funnel is folded over, resulting in a quid, which is placed in the mouth and chewed.^{2,9}

Chemical Measurements

Published chemical data for betel quid are not available.

References

1. Reddy KS, Gupta PC, editors. Report on tobacco control in India. New Delhi: Ministry of Health and Family Welfare, Government of India; 2004. Available from: http://www.who.int/fctc/reporting/Annex6_Report_on_Tobacco_Control_in_India_2004.pdf
2. International Agency for Research on Cancer. Betel-quid and areca-nut chewing and some areca-nut-derived nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 85. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2004. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol85/mono85.pdf>
3. The National. UAE police close chewing tobacco factories to clean up towns and cities. The National. 2009 March 31 [cited 2012 Oct 16]. Available from: http://www.dajjworld.com/news/news_disp.asp?n_id=58384&n_fit=UAE%3A+Police+Close+Chewing+Tobacco+Factories+to+Clean+Up+Towns+%26+Cities
4. World Health Organization, Regional Office for the Western Pacific. Review of areca (betel) nut and tobacco use in the Pacific: a technical report. Manila: World Health Organization, Regional Office of the Western Pacific; 2012 [cited 2012 Aug 16]. Available from: <http://www.wpro.who.int/tobacco/documents/betelnut.pdf>
5. International Institute for Population Sciences, Ministry of Health and Family Welfare, Government of India. Global Adult Tobacco Survey, India, 2009–2010. New Delhi: Ministry of Health and Family Welfare; Mumbai: International Institute for Population Sciences; 2010 [cited 21 June 2012]. Available from: http://whoindia.org/EN/Section20/Section25_1861.htm
6. Ministry of Health and Family Welfare (Bangladesh). Global Adult Tobacco Survey: Bangladesh report 2009. Dhaka, Bangladesh: World Health Organization, Country Office for Bangladesh; 2009. Available from: http://www.who.int/tobacco/surveillance/global_adult_tobacco_survey_bangladesh_report_2009.pdf
7. Lee CH, Ko AM, Warnakulasuriya S, Yin BL, Sunarjo, Zain RB, et al. Intercountry prevalences and practices of betel-quid use in south, southeast and eastern Asia regions and associated oral preneoplastic disorders: an international collaborative study by Asian Betel-Quid Consortium of South and East Asia. *Int J Cancer*. 2011;129(7):1741–51.
8. World Health Organization. Myanmar sentinel tobacco use prevalence study, 2004. World Health Organization; [no year]. Available from: http://209.61.208.233/LinkFiles/Regional_Tobacco_Surveillance_System_Myanmar2004.pdf
9. Gupta PC, Ray CS. Epidemiology of betel quid usage. *Ann Acad Med Singapore*. 2004;33(4 Suppl.):31–6.
10. Reichart PA, Nguyen XH. Betel quid chewing, oral cancer and other oral mucosal diseases in Vietnam: a review. *J Oral Pathol Med*. 2008 Oct;37(9):511–4. Epub 2008 Jul 8.

For additional information on betel quid (paan), please refer to the following chapters: Chapter 10: Smokeless Tobacco Use in the European Region; Chapter 11: Smokeless Tobacco Use in the Eastern Mediterranean Region; Chapter 13: Smokeless Tobacco Use in the South-East Asia Region; and Chapter 14: Smokeless Tobacco Use in the Western Pacific Region.

Chimó

Chimó is a tobacco-based paste that was reportedly first used in South America in the early days of European colonization. In 1497, Amerigo Vespucci reported the use of a chewing tobacco mixed with ashes in the Caribbean.¹ According to a popular legend, the aboriginal chief Chimauchu was the first to use tobacco in the form of a paste, which today is called chimó.¹



Photo courtesy of Clifford Watson,
Centers for Disease Control and Prevention

Common Names

None

Brand Names

El Tovareño, El Tigrito, El Sabroso, El Gran Búfalo, El Dragón, El Morichal, San Carleño

Main Geographic Locations

(WHO Region: Country)

Region of the Americas: Venezuela, Colombia²

Prevalence and Demographics

Among adults in Venezuela in 2007, 1.5% of women and 6.2% of men were current users, and 3.1% of women and 15.4% of men reported ever using chimó.³ The Venezuelan Global Youth Tobacco Survey (GYTS) was used to estimate tobacco-specific prevalence of smokeless use among students in grades 7–9. GYTS results for Venezuela nationally and for eight different states in the years 2000, 2004, and 2008 found that the prevalence of chimó use was not uniform among the states: It ranged from 3.8% to 20.7% for boys, and 2.0% to 6.6% for girls.² Prevalence rates of chimó use in Colombia are not available.

Mode of Absorption

Oral (sucked, held in mouth)

Use Pattern

A small amount of chimó is placed between the lip or cheek and the gum and left there for some time, usually 30 minutes. The mixture of chimó and saliva is spit out.⁴

Main Ingredients

Tobacco leaf, baking soda (sodium bicarbonate), brown sugar, ashes from the Mamón tree (*Melicocca bijuga*), and vanilla and anisette flavoring. The ingredients vary according to the region.⁴

Processing/Manufacturing

Cottage industry and commercial: Chimó is usually produced by small family-operated factories, but commercial, industrial manufacturing of chimó is increasing in Venezuela. Tobacco leaves are first crushed and boiled for several hours or days, and then starch and fibers are removed. The remaining concentrated product (10 kilos of tobacco becomes one kilo of “basic” chimó paste) is a sticky, heavy, black liquid, which can be stored for maturation for up to 2 years. For maturation, it is placed in natural containers like “taparas” (dried fruit from the Tapara tree) or wrapped in banana leaves. The matured paste is seasoned with other ingredients (see “Main Ingredients”), and then packaged in small tins or candy-like wrapped cylinders.⁴

Chimó

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNAs*
		mg/g wet wt		ng/g wet wt			
Chimó, Venezuela	6.98–9.40	5.29–30.1	1.32–27.4	310–2,600	318–4,260	14.9–1,330	954–9,390

*Total TSNAs represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).
Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosonornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAs = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.
Source: Stanfill et al. 2011 (5).

References

1. Monteverde HR, Magaña AR. Breves comentarios sobre la historia del tabaco y el tabaquismo. *Rev Inst Nal Enf Resp Mex.* 2006;19(4):297–300. Available from: http://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S0187-75852006000400013&lng=es. Spanish.
2. Granero R, Jarpa P. Uso de chimo entre adolescentes en Venezuela. *Encuesta Mundial sobre Tabaquismo en Jóvenes 1999–2008. Acta Odontol Venezolana.* 2011;49(3). Spanish.
3. Infante E, Navarro A, Finizola R, Zevallos J, Moya D, Alvarado O, et al. Estudio de la salud cardiovascular en el Estado Lara (ESCEL 2008) metodología del proyecto y perfil de la muestra estudia. *Avances Cardiol.* 2009;30(4):316–25. Spanish.
4. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
5. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. *Tob Control.* 2011 May;20(3):e2. Epub 2010 Nov 25.

For additional information on chimó products, please refer to Chapter 9: Smokeless Tobacco Use in the Region of the Americas.

Creamy Snuff

Creamy snuff is a tobacco-based paste sold in toothpaste-like tubes. It is often advertised as being antibacterial and healthy for teeth and gums, and it is used primarily by women.¹



Photo courtesy of World Health Organization South-East Asia Regional Office and Dharendra N. Sinha

Common Names

Tobacco toothpaste

Brand Names

IPCO (Asha Industries), Dentobac (Goran Pharma LTD), Tona, Ganesh, Charotar, Musa Ka, Rehmat Khan, Chad Tara, Dulhan, Suraj, Asif Ka

Main Geographic Location

(WHO Region: Country)

South-East Asia Region: India²

Prevalence and Demographics

Creamy snuff is used primarily by women, but it also seems popular among children.^{1,3} In 2004, one study in India reported that the prevalence of creamy snuff use among adolescents aged 13 to 15 years varied from 2% to 32% across 18 states.⁴ Although no product-specific adult prevalence data are available, in 2009–2010 4.7% of all adults (3.3% of males and 6.3% of females) in India reported applying tobacco products, including creamy snuff, mishri, gul, or gudakhu, to their teeth and gums.⁵

Mode of Absorption

Oral (applied to teeth and gums, teeth cleaning)

Use Pattern

Creamy snuff is used to clean teeth like regular toothpaste. Some products' instructions recommend holding the paste in the mouth for a little while before rinsing.^{1,3}

Main Ingredients

Tobacco, clove oil, glycerin, spearmint, menthol, camphor, water³

Processing/Manufacturing

Commercial: Creamy snuff is commercially manufactured and is marketed as a teeth cleaner (dentifrice).³ Creamy snuff consists of finely ground tobacco mixed with aromatic substances, such as clove oil, glycerin, spearmint, menthol and camphor, salts, water, and other hydrating agents.^{1,3} Additional information on the manufacturing of creamy snuff could not be located.

Creamy Snuff

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type*	pH	Total Nicotine mg/g wet wt	Free Nicotine mg/g wet wt	NNK	NNN	NNAL	Total TSNAs†
Creamy Snuff, India	7.51–8.35	5.62–10.0	2.36–3.82	N/A	N/A	N/A	N/A

*All creamy snuff products were commercially manufactured.

†Total TSNAs represent the sum of NNK, NNN, and NNAL (shown), *N*'-nitrosoanatabine and *N*'-nitrosoanabasine (not shown).

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N*'-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAs = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Gupta and Sreevidya 2004 (6).

References

1. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf
2. Gupta PC, Ray CS. Smokeless tobacco and health in India and South Asia. *Respirology*. 2003;8(4):419–31.
3. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
4. Sinha DN, Gupta PC, Pednekar MS. Use of tobacco products as dentifrice among adolescents in India: questionnaire study. *Brit Med J*. 2004;328(7435):323–4.
5. International Institute for Population Sciences, Ministry of Health and Family Welfare, Government of India. Global Adult Tobacco Survey, India, 2009–2010. New Delhi: Ministry of Health and Family Welfare; Mumbai: International Institute for Population Sciences; 2010 [cited 21 June 2012]. Available from: http://whoindia.org/EN/Section20/Section25_1861.htm
6. Gupta PC, Sreevidya S. Report on smokeless tobacco products in India. Mumbai: Action Council Against Tobacco; 2004.

Dissolvable Tobacco

Dissolvable smokeless tobacco products were introduced in the United States in 2001. The first products, Ariva and Stonewall, dissolved completely during use, with no residual loose tobacco or tobacco-containing sachet to discard. In 2009, R.J. Reynolds Tobacco Company released Camel Orbs, Strips, and Sticks in a variety of formulations and flavors, all of which can fully dissolve in the mouth.¹ In 2011, Altria began test marketing new dissolvable products, Marlboro and Skoal Smokeless Tobacco Sticks, which are small toothpick-like sticks coated with tobacco, which are discarded after the coating dissolves.² In January 2013, the manufacture and sale of Ariva and Stonewall lozenges were discontinued.³ Because dissolvables look like candy, there is concern that children will accidentally ingest them.⁴



Photo courtesy of Clifford Watson, Centers for Disease Control and Prevention

Type	Description	Examples
Tablets	Tobacco compressed into a lozenge	Ariva, Camel Orbs, Stonewall
Strips	Tobacco pressed into a thin wafer-like dissolvable strip (somewhat like a breath strip)	Camel Strips
Sticks	Tobacco pressed into a long, thin dissolvable stick	Camel Sticks
Tobacco Sticks	Small toothpick-like stick coated with a semi-hard tobacco-containing mixture. All of this product dissolves in the mouth except the toothpick.	Marlboro Tobacco Sticks, Skoal Tobacco Sticks

Common Names

Dissolvables

Brand Names

Ariva, Stonewall (*Star Scientific*); Camel Orbs, Camel Strips, Camel Sticks (*R.J. Reynolds*); Marlboro Sticks (*Philip Morris*); Skoal Sticks (*U.S. Smokeless Tobacco Company*)

Main Geographic Location (WHO Region: Country)

Region of the Americas: United States⁵

Prevalence and Demographics

A 2010 survey found that less than 1% (0.6%) of adults in the United States had tried dissolvables.⁶ Dissolvable tobacco products are fairly new to the market, and as of 2011–2012, Camel, Marlboro,

and Skoal dissolvable products were still being test marketed in select cities and were not nationally available.^{2,5,7} As these products become more widely available, their use may become more widespread.

Mode of Absorption

Oral (sucked, held in mouth, dissolved)

Use Pattern

These products can dissolve in as few as 3 minutes (Camel Orbs) or as much as 30–60 minutes (Stonewall).⁸ Tobacco sticks do not completely dissolve because the tobacco is coated onto a wooden toothpick that must be thrown away.

Main Ingredients

Tobacco, humectants, preservatives, flavors⁹

Dissolvable Tobacco

Processing/Manufacturing

Commercial: Dissolvable tobacco is commercially manufactured. Limited information is available on the specific processes used for manufacturing many of these products. To create tobacco tablets,

tobacco is pasteurized and ground into fine powder, and the finely powdered tobacco is combined with flavors, such as mint and eucalyptus, and other additives that allow the tobacco to be compressed into tablet form and dissolved in the mouth.¹⁰

Chemical Measurements

These data are for select products and may not represent all products of this type. Data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type*	pH	Total Nicotine mg/g wet wt	Free Nicotine mg/g wet wt	NNK	NNN	NNAL	Total TSNAs†
Tablets,‡ United States	6.92–8.10	3.03–6.85	0.37–1.65	69–280	61–214	—	536–736
Camel Sticks, United States	7.76	3.92	1.45	307	264	—	852
Camel Strips, United States	7.88	2.67	1.11	221	152	—	535

*All products were commercially manufactured.

†Total TSNAs represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).

‡Includes Camel Orbs, Ariva, and Stonewall.

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosonornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAs = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Stepanov et al. 2012 (11).

References

1. R.J. Reynolds Tobacco Company. Products at a glance: Camel Sticks, Strips and Orbs. 2008. Available from: http://www.ftac.org/tcn/ftp/2008/october-2008/pdf/RJR_Camel_Dissolvables_info.pdf
2. Abu-Shalback Zid L. Altria Smokeless Tobacco Sticks. Tobacco E-News [Internet]. 2011 Apr 5 [cited 2012 Sept 24]. Available from: <http://www.cspnet.com/news/tobacco/articles/altria-smokeless-tobacco-sticks>
3. Star Scientific, Inc. Subsidiaries & products: Star Tobacco, Inc. 2012 Dec [cited 2013 Jan 28]. Available from: <http://www.starscientific.com/products/star-tobacco-inc/>
4. Connolly G, Richter P, Aleguas A, Stanfill S, Alpert H. Unintentional child poisoning through ingestion of conventional and novel tobacco products. *Pediatrics*. 2010;125(5):896–9. Epub 2010 Apr 19.
5. Mann D. FDA to weigh safety of tobacco lozenges, strips. *USA Today*. 2012 Jan 18 [cited 2012 Sept 24]. Available from: <http://www.usatoday.com/news/health/story/health/story/2012-01-18/FDA-to-weigh-safety-of-tobacco-lozenges-strips/52631210/1>
6. McMillen R, Maduka J, Winickoff J. Use of emerging tobacco products in the United States. *J Environ Public Health*. 2012; Article ID 989474. Epub 2012 May 10.
7. The Alliance for Safe and Drug Free Children. Camel Sticks, Strips & Orbs – Alert. Alliance Newsletter. 2011 July 7 [cited 2012 Oct 2]. Available from: <http://www.drug-alliance.org/newsletter/camel-sticks-strips-orbs-alert>
8. Varney SM. Dissolvable tobacco products. ACEP Toxicology Section Newsletter. 2009;16(1) [cited 2012 Sept 24]. Available by subscription from: www.acep.org
9. Rainey C, Conder P, Goodpaster J. Chemical characterization of dissolvable tobacco products promoted to reduce harm. *J Agr Food Chem*. 2011;59(6):2745–51.
10. Philip Morris Products SA. Smokeless dissolvable compressed tobacco product [patent application]. Pub No WO/2010/086010. World Intellectual Property Organization. 2010 [cited 2012 Sept 24]. Available from: <http://patentscope.wipo.int/search/en/detail.jsf?docId=WO2010086010&recNum=1&maxRec=&offce=&prevFilter=&sortOption=&queryString=&tab=PCT+Biblio>
11. Stepanov I, Biener L, Knezevich A, Nyman AL, Bliss R, Jensen J, et al. Monitoring tobacco-specific *N*-nitrosamines and nicotine in novel Marlboro and Camel smokeless tobacco products: findings from Round 1 of the New Product Watch. *Nicotine Tob Res*. 2012 Mar;14(3):274–81. Epub 2011 Oct 29.

For additional information on dissolvables, please refer to Chapter 9: Smokeless Tobacco Use in the Region of the Americas.

Dohra



Photo courtesy of World Health Organization South-East Asia Regional Office and Dharendra N. Singh

Dohra is a wet mixture of tobacco and other ingredients such as areca nut, catechu, and flavors. Tobacco is frequently added to this mixture in the form of zarda, a smokeless tobacco product composed of flavored tobacco flakes. Dohra is commonly used in the Allahabad District of the Uttar Pradesh state in India, and in surrounding districts, such as Jaunpur and Pratapgarh.¹

Common Names

None

Brand Names

None

Main Geographic Location

(WHO Region: Country)

South-East Asia Region: India (Uttar Pradesh)²

Prevalence and Demographics

It is estimated that more than 50% of people in the Jaunpur area of India use dohra.² There are no available data on nationally representative prevalence rates of dohra use in India.

Mode of Absorption

Oral (chewed)

Use Pattern

Users may buy the dohra spice mixture packet and tobacco packet separately and add the amount of tobacco they prefer.^{1,2}

Main Ingredients

Tobacco, areca nut, and other ingredients such as catechu, slaked lime (calcium hydroxide), peppermint, cardamom^{1,2}

Processing/Manufacturing

Custom-made: Dohra is produced by individual vendors for sale. It is sold either as a ready-made mixed tobacco product or in two packets, one containing tobacco (often zarda or surti) that is mixed with the contents of the second packet (areca nut, catechu, and other flavorings). Dohra is normally sold in a plastic bag with a rubber band tied around it.²

Chemical Measurements

No data available

References

1. Pandya S, Chaudhary A, Singh M, Singh M, Mehrotra R. Correlation of histopathological diagnosis with habits and clinical findings in oral submucous fibrosis. *Head Neck Oncol.* 2009;1:10. Epub 2009 May 2.
2. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/enfity/tobacco/topics/oral_tobacco_use.pdf

Dry Snuff

Two types of snuff are manufactured and used in the United States: moist snuff and dry (Scotch) snuff. Moist snuff is by far the most widely consumed type in the United States and Canada. It contains 20%–60% moisture and often is flavored with wintergreen or various fruit flavors. In contrast, dry snuff, a finely powdered tobacco product produced mainly from Kentucky and Tennessee fire-cured tobaccos, has a moisture content that is less than 10% by weight.¹ This factsheet describes only commercial dry snuff. Other types of dry snuff are used traditionally in many regions around the world (see the factsheets *Tapkheer* and *African Traditional Snuff Products* for more information).



Photo courtesy of Clifford Watson, Centers for Disease Control and Prevention

Common Names

Scotch snuff, snuff

Brand Names

Levi Garrett & Sons, Dental, Honest, Peach Sweet, Tube Rose, W.E. Garrett & Sons (*American Snuff Company*); Silver Dollar (*Kreteck International, Inc.*)

Main Geographic Locations

(WHO Region: Country)

Region of the Americas: Canada, United States¹;

African Region: South Africa, Nigeria²;

European Region: Germany¹

Prevalence and Demographics

Use of dry snuff has declined over the last 100 years and is rare today. In 2012, the U.S. prevalence of smokeless tobacco use (including snuff and chewing tobacco) among those aged 12 years and older was 3.6% (7.1% of males and 0.4% of females).³ Although no current prevalence data are available specifically for dry snuff use, sales of dry snuff

comprised only 1.4% of all smokeless tobacco sales in the United States in 2009.⁴ Prevalence data on dry snuff use in other countries are not available.

Mode of Absorption

Oral (sucked, held in mouth); **Nasal**

Use Pattern

Dry snuff is usually used orally, but it may also be inhaled into the nostrils.¹

Main Ingredients

Tobacco (fire-cured, fermented), often flavored¹

Processing/Manufacturing

Commercial: Dry snuff is commercially manufactured. Tobacco is fire-cured and then fermented and processed into a dry, powdered form; it may also be sweetened. The moisture content of the finished product is less than 10% by weight. It is packaged and sold in small metal or glass containers.¹

Dry Snuff

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type*	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNAst
		mg/g wet wt		ng/g wet wt			
Dry Snuff (5), United States	5.71–6.25	14.9–20.2	0.07–0.30	1,340–14,600	6,120–31,300	47–1,050	10,300–76,500
Dry Snuff (1), United States	5.41–7.96	4.70–24.84	0.03–3.13	—	—	—	—

*Dry snuff products were commercially manufactured.

†Total TSNAst represent the sum of NNK, NNN, and NNAL (shown), *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAst = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Sources: Lawler et al. 2013 (5); International Agency for Research on Cancer 2007, table 6 (1).

References

1. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
2. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. *Tob Control*. 2011 May;20(3):e2. Epub 2010 Nov 25.
3. Substance Abuse and Mental Health Services Administration. Results from the 2012 National Survey on Drug Use and Health: detailed tables. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2013. Available from: <http://www.samhsa.gov/data/NSDUH/2012SummNatFindDetTables/Deftabs/NSDUH-DeftabsLOTsect2pe2012.htm#TopOfPage>
4. Maxwell J. The Maxwell report: the smokeless tobacco industry in 2009. Richmond, VA: John C. Maxwell, Jr.; 2010.
5. Lawler TS, Stanfill SB, Zhang L, Ashley DL, Watson CH. Chemical characterization of domestic oral tobacco products: total nicotine, pH, unprotonated nicotine and tobacco-specific *N*-nitrosamines. *Food Chem Toxicol*. 2013;57:380-6. Epub 2013 Mar 19. doi 10.1016/j.fct.2013.03.011

For additional information on dry snuff, please refer to the following chapters: Chapter 9: Smokeless Tobacco Use in the Region of the Americas, and Chapter 12: Smokeless Tobacco Use in the African Region.

Gudakhu

Gudakhu (also spelled gudahku) is a paste-like product that is made from fine tobacco leaf dust and molasses (called *sheera*), red soil, and lime.^{1,2} Gudakhu is predominantly used in India in the states of Bihar, Orissa, Uttar Pradesh, and Uttaranchal.² It may be used to clean teeth.^{3,4}



Photo courtesy of World Health Organization South-East Asia Regional Office and Dharendra N. Sinha

Common Names

None

Brand Names

Natraj

Main Geographic Location

(WHO Region: Country)

South-East Asia Region: India⁴

Prevalence and Demographics

No recent information was located specifically on the use of gudakhu. Gudakhu is mainly used by women. Survey data from the 1970s found that 1% of men and 16% of women in eastern India (Jharkhand) used gudakhu.^{2,4} In 2009–2010, 4.7% of all adults (3.3% of males and 6.3% of females) in India reported use of tobacco products that are applied to the teeth and gums, including gudakhu, snuff, mishri, or gul.⁵

Mode of Absorption

Oral (applied to teeth and gums, held in mouth)

Use Pattern

Gudakhu can be rubbed on the teeth and gums with a fingertip and may be left in the mouth. It is an addictive substance, with some people using it up to 20 times per day.¹

Main Ingredients

Tobacco powder, molasses, red soil, lime, water^{1,4}

Processing/Manufacturing

Commercial and custom-made: Gudakhu is available commercially, but can also be made by individuals for personal use. It can come in different types of packaging (both branded and unbranded) and is frequently carried in a metal container.² Additional information on manufacturing of gudakhu could not be located.

Chemical Measurements

No data available

References

1. Dash BC, Das RK. Genotoxicity of 'gudakhu', a tobacco preparation. I. In mice in vivo. *Mutat Res.* 1992;280(1):45–53.
2. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf
3. National Cancer Institute. Smokeless tobacco or health: an international perspective. Smoking and tobacco control monograph no. 2. Bethesda, MD: U.S. Department of Health and Human Services, National Cancer Institute; 1992. NIH publication no. 93-3461. Available from: <http://www.cancercontrol.cancer.gov/tcrb/monographs/2/index.html>
4. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific N-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
5. International Institute for Population Sciences, Ministry of Health and Family Welfare, Government of India. Global Adult Tobacco Survey, India, 2009–2010. New Delhi: Ministry of Health and Family Welfare; Mumbai: International Institute for Population Sciences; 2010 [cited 21 June 2012]. Available from: http://whoindia.org/EN/Section20/Section25_1861.htm

Gul

Gul is a pyrolysed (burned and decomposed), powdered tobacco product that is marketed in small tin cans or sachets under several different brand names. It is used as a dentifrice in India.¹



*Photo courtesy of Clifford Watson,
Centers for Disease Control and Prevention*

Common Names

None

Brand Names

Shajadi Gul, Mujamal Hussain Musarraf Bahi Shahi Eagle, Md. Mustafa Asgar Ali Gul (Bangladesh), Chand, Tara Marka, and Gulbadan (India)

Main Geographic Locations (WHO Region: Country)

South-East Asia Region: India, Bangladesh¹

Prevalence and Demographics

In 2009, 5.3% of adults (5.5% of males and 5.1% of females) in Bangladesh reported using gul.² In India, gul is popular among women,^{3,4} and in 2004, 2–6% of adolescents in various regions of India reported using it.⁵ There are no recent nationally representative statistics on the prevalence of gul use in India. In 2009–2010, 4.7% of all adults in India (3.3% of males and 6.3% of females) reported using oral tobacco, including gul, snuff, mishri, or gudakhu.⁴

Mode of Absorption

Oral (teeth cleaning, applied to teeth and gums)

Use Patterns

Gul is usually used to clean teeth. It is addictive and may be used several times a day.^{1,3}

Main Ingredients

Pyrolysed (burned and decomposed) tobacco leaves, molasses, other unknown ingredients^{3,6}

Processing/Manufacturing

Commercial: Gul is commercially manufactured³ and sold in small tin cans.⁶ Additional information on manufacturing of gul could not be located.

Gul

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type*	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNAst
		mg/g wet wt					
Gul Powder, Bangladesh	8.79–9.22	33.4–34.1	29.1–31.0	1,330–1,370	5,190–8,020	590–630	13,400–17,100

*Gul products were commercially manufactured.

†Total TSNAst represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown). Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrososornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAst = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Stanfill et al. 2011 (7).

References

1. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf
2. Ministry of Health and Family Welfare (Bangladesh). Global Adult Tobacco Survey: Bangladesh report. Dhaka, Bangladesh: World Health Organization, Country Office for Bangladesh; 2009. Available from: http://www.who.int/tobacco/surveillance/global_adult_tobacco_survey_bangladesh_report_2009.pdf
3. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
4. International Institute for Population Sciences, Ministry of Health and Family Welfare, Government of India. Global Adult Tobacco Survey, India, 2009–2010. New Delhi: Ministry of Health and Family Welfare; Mumbai: International Institute for Population Sciences; 2010 [cited 21 June 2012]. Available from: http://whoindia.org/EN/Section20/Section25_1861.htm
5. Sinha DN, Gupta PC, Pednekar MS. Use of tobacco products as dentifrice among adolescents in India: questionnaire study. *Brit Med J*. 2004;328(7435):323–4.
6. Reddy KS, Gupta PC, editors. Report on tobacco control in India. New Delhi: Ministry of Health and Family Welfare, Government of India; 2004. Available from: http://www.who.int/fctc/reporting/Annex6_Report_on_Tobacco_Control_in_India_2004.pdf
7. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. *Tob Control*. 2011 May;20(3):e2. Epub 2010 Nov 25.

For additional information on gul, please refer to the following chapters: Chapter 13: Smokeless Tobacco Use in the South-East Asia Region.

Gutka

(Pan Masala with Tobacco)

Gutka, or gutkha, is a dry, granular tobacco that is a commercially manufactured version of pan or betel quid with tobacco.¹ Unlike pan, gutka is relatively nonperishable and packaged in small single-use packets that are easy to carry and use “anytime, anywhere.”² Advertising of tobacco products on radio and TV has been banned in India since 2004; however, pan masala, an identical product without tobacco but the same brand names, can be advertised, and these ads are often targeted at youth.¹ As of October 2013, almost all states and union territories in India have banned gutka, although it is unclear how well these bans will be enforced.^{3,4}



Photo courtesy of World Health Organization South-East Asia Regional Office and Dharendra N. Sinha

Common Names

Pan masala. (The terms gutka and pan masala are sometimes used interchangeably, but gutka usually refers to the product with tobacco, and pan masala is the same product without tobacco.⁵)

Brand Names

Manikchand, Moolchand, Tulsi, Shimla, Sikandar, Pan Parag, RMD, Sir, Shikhar, Dandia, Kuber, Wiz, Kesar, Club Class, Goa, Shanti Strong, Vimal, Zee, Mehak Silver, Silver, Kanchan

Packaging of some brands of gutka is often identical to packaging of pan masala (which does not contain tobacco), and companies may be using this identical packaging to circumvent India's 2004 ban on tobacco advertising.¹

Main Geographic Locations

(WHO Region: Country)

South-East Asia Region: India, Bangladesh, Nepal, Myanmar, Sri Lanka^{2,6}; **Eastern Mediterranean Region:** Pakistan⁵

Prevalence and Demographics

In 2009–2010, 8.2% of all individuals in India aged 15 years and older (13.1% of males and 2.9% of females) reported chewing gutka.⁵ Use of gutka is common among youth and young adults.^{1,5,7,8} A number of surveys conducted in India have shown

that pan masala and gutka are commonly chewed by children and adolescents.^{5,9} Product-specific prevalence data are not available for the other countries where gutka is used.

Mode of Absorption

Oral (chewed, held in mouth)

Use Patterns

Gutka is held in the mouth and chewed. Saliva is generally spit out, but it is also sometimes swallowed.⁶ These products are commonly used throughout the day.

Main Ingredients

Tobacco, areca nut, slaked lime (calcium hydroxide), catechu, and other condiments, sweeteners, and flavorings⁶

Processing/Manufacturing

Commercial and cottage industry: Gutka is generally commercially manufactured as a dry, relatively nonperishable preparation.¹ Gutka can also be a premade cottage product¹⁰ that is packaged in nontraditional packaging (i.e., cellophane). Gutka is made from powdered tobacco, areca nut, lime, and catechu, with other condiments and sweeteners added for flavor.² Manufactured gutka is sold in small, brightly colored plastic and paper packets (sachets); plastic sachets were banned in India in March 2011.¹¹

Gutka

(Pan Masala with Tobacco)

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNA ^s *
		mg/g wet wt		ng/g wet wt			
Gutka,† Pakistan	8.20–8.54	0.16–2.08	0.12–1.08	11.6–208	45.4–913	7.02–53.5	83.9–1,560
Gutka,‡ India	8.46–8.88	1.09–2.33	0.86–1.78	57.1–456	167–1,280	23.2–258	370–2,250
Gutka,§ India	7.43–8.61	0.91–4.20	0.19–3.33	7.1–375	154–18,600	10.8–1,030	264–23,900

*Total TSNA^s represent the sum of NNK, NNN, and NNAL (shown) and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).

†These gutka products were manufactured both commercially and by cottage industry.

‡These gutka products were commercially manufactured.

§These gutka products were manufactured by cottage industry.

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNA^s = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Stanfill et al. 2011 (10).

References

- Gupta PC, Ray CS. Epidemiology of betel quid usage. *Ann Acad of Med Singapore*. 2004;33(4 Suppl):31–6.
- World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf
- Arora M, Madhu R. Banning smokeless tobacco in India: policy analysis. *Indian J Cancer*. 2012;49(4):336–41.
- Economic Times Bureau. Health Ministry endorses ban on chewing tobacco, gutka. *The Economic Times*; 2013 Oct 2 [cited 2013 Oct 29]. Available from: http://articles.economicstimes.indiatimes.com/2013-10-02/news/42617601_1_smokeless-tobacco-indirect-morbidity-costs-tobacco-use
- International Agency for Research on Cancer. Betel-quid and areca-nut chewing and some areca-nut-derived nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 85. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2004. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol85/mono85.pdf>
- International Agency for Research on Cancer. Smokeless tobacco and some tobacco-related *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
- International Institute for Population Sciences, Ministry of Health and Family Welfare, Government of India. Global Adult Tobacco Survey, India, 2009–2010. New Delhi: Ministry of Health and Family Welfare; Mumbai: International Institute for Population Sciences; 2010 [cited 21 June 2012]. Available from: http://whoindia.org/EN/Section20/Section25_1861.htm
- Gupta PC, Ray CS. Smokeless tobacco and health in India and South Asia. *Respirology*. 2003;8(4):419–31.
- Narain R, Sardana S, Gupta S, Sehgal A. Age at initiation and prevalence of tobacco use among school children in Noida, India: a cross-sectional questionnaire based survey. *India J Med Res*. 2011;133(3):300–7.
- Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. *Tob Control*. 2011 May;20(3):e2. Epub 2010 Nov 25.
- Singh N, Zachariah R. Plastic ban to hit sachet makers. *The Times of India*; 2011 Feb 9 [cited 2 Oct 2012]. Available from: http://articles.timesofindia.indiatimes.com/2011-02-09/india-business/28543748_1_pan-masala-gutka-plastic-ban

For additional information on gutka, please refer to the following chapters: Chapter 10: Smokeless Tobacco Use in the European Region, and Chapter 13: Smokeless Tobacco Use in the South-East Asia Region.

Iqmik

Iqmik is a homemade chewing tobacco that consists of tobacco leaves combined with ash from burnt tree fungus or wood.^{1,2,3} Iqmik is commonly used by Alaska Native people, particularly the Yup'ik and Cup'ik Eskimo people in western Alaska.^{2,4} In contrast to American Indian customs, tobacco use does not have a spiritual significance for Alaska Natives.^{1,2} Rather, it is a relatively recent phenomenon, dating back only about 150 years in Alaska, although tobacco use among other indigenous populations in North America dates back at least 2,000 years.^{2,3} Some people in the Yukon–Kuskokwim region consider iqmik a healthier alternative to smoking tobacco because its ingredients are perceived as “natural.”^{1,2}



*Photo courtesy of Caroline Renner,
Alaska Native Medical Center*

Common Names

Blackbull, dediguss

Brand Names

None

Main Geographic Location (WHO Region: Country)

Region of the Americas: United States (Alaska)⁵

Prevalence and Demographics

Iqmik is used by Alaska Natives, including youth of all ages.^{1,2} Iqmik is used in western Alaska, although there are reports of similar smokeless tobacco mixtures being used by indigenous populations in Western Siberia, Yukon, Labrador, British Columbia Coast, and Nova Scotia.^{2,4} Data from the 2004 to 2007 Behavioral Risk Factor Surveillance System (BRFSS) found that iqmik is used by 16% to 22% of Alaska Native adults in the western region of the state.^{6,7} It has also been reported that iqmik is often used by pregnant women in western Alaska.⁴

Mode of Absorption

Oral (chewed)

Use Pattern

Users may pre-chew the iqmik and place it in a small box for later use by the maker or to share with others, including elders, children, and infants. Iqmik is believed by some to relieve babies' teething pain.^{1,5}

Main Ingredients

Tobacco, tree fungus ash (also known as punk, araq, or buluq ash) or other ash derived from burning driftwood, alder bush, or willow bush; these woods are used because there are few trees along the coast of western Alaska.^{2,4}

Processing/Manufacturing

Custom-made: Although the ingredients are available at grocery stores and retail outlets, iqmik is always prepared by the individual user or a family or community member to share. Fire- or air-cured tobacco leaves are mixed with punk fungus ash, which is generated by burning a woody fungus that grows on birch trees, or other woody ash if punk ash is unavailable.^{1,2,5} The mixture can be prepared in several ways: by pre-chewing, stirring in a bowl with water, or even using a modern blender.^{1,4} Iqmik is often stored in a commercial smokeless tobacco box or a small plastic container.^{2,4}

Iqmik

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for the prepared iqmik. The amount absorbed will depend on the amount of product used.

Product Type	pH*	Total Nicotine Free Nicotine mg/g wet wt		NNK	NNN	NNAL	Total TSNAst
Iqmik with Air-Cured Tobacco, ‡ United States (Alaska)	11.0	38.3	38.3	209	2,995	61	7,238
Iqmik with Fire-Cured Tobacco, ‡ United States (Alaska)	11.0	38.9	38.9	473	2,400	11	7,191

*pH of the prepared ash/tobacco iqmik mixture; both willow and punk ash are pH 11.0.

†Total TSNAst represent the sum of NNK, NNN, and NNAL (shown), and *N*'-nitrosoanatabine and *N*'-nitrosoanabasine (not shown).

‡Cured tobacco includes the average of twist and leaf tobacco.

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N*'-nitrosonornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAst = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Hearn et al. 2013 (8).

References

1. Renner C, Patten C, Enoch C, Petraitis J, Offord K, Angstman S, et al. Focus group of Y-K Delta Alaska Natives: attitudes toward tobacco use and tobacco dependence interventions. *Prev Med.* 2004;38(4):421–31.
2. Blanchette R, Renner C, Held B, Enoch C, Angstman S. The current use of *Phellinus igniarius* by the Eskimos of western Alaska. *Mycologist.* 2002 Nov;16(4):142–5.
3. Wolsko C, Mohatt GV, Lardon C, Burket R. Smoking, chewing, and cultural identity: prevalence and correlates of tobacco use among the Yup'ik: the Center for Alaska Native Health Research (CANHR) study. *Cult Divers Ethn Min Psychol.* 2009;15(2):165–72.
4. Renner CC, Enoch C, Patten CA, Ebbert JO, Hurt RD, Moyer TP, et al. Iqmik: a form of smokeless tobacco used among Alaska Natives. *Am J Health Behav.* 2005;29(6):588–94.
5. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
6. Alaska Behavioral Risk Factor Surveillance System (BRFSS) 2004 [data]. Anchorage: Alaska Department of Health and Social Services; 2005.
7. Alaska Behavioral Risk Factor Surveillance System (BRFSS) 2007 [data]. Anchorage: Alaska Department of Health and Social Services; 2007.
8. Hearn BA, Renner C, Vaughan-Watson C, Stanfill SB, Zhang L, et al. Chemical analysis of Alaskan iq'mik smokeless tobacco. *Nic Tob Res.* 2013;15(7):1283-8. doi: 10.1093/ntr/nts270

For additional information on iqmik, please refer to Chapter 9: Smokeless Tobacco Use in the Region of the Americas.

Khaini

Khaini (or khoinee in Bangladesh) is a flaky product made of sun-dried tobacco and slaked lime. Lime is often added to the tobacco just before use.¹ Khaini is used in India, Nepal, and Bangladesh.²



Photo courtesy of Clifford Watson,
Centers for Disease Control and Prevention

Common Names

Chada, chadha, sada, surti (in Nepal and neighboring parts of India)

Brand Names

Raja, Kuber, Wiz, Buddha Lal, Chaini, Raja Chap, Ansul Tobacco, Mirage, Ganesh Tobacco 701, Patta Chhap Tej Tobacco

Main Geographic Locations (WHO Region: Country)

South-East Asia Region: India, Bangladesh, Nepal, Bhutan²

Prevalence and Demographics

Among adults in India, khaini is the most commonly used type of smokeless tobacco. In 2009–2010, 11.6% of the Indian population over age 15 (18% of males and 4.7% of females) used khaini.³ In 2009, in Bangladesh, 1.5% of all adults (1.9% of males and 1.2% of females) used khaini.⁴ Product-specific prevalence rates were not available for the other countries where khaini is used.

Mode of Absorption

Oral (sucked, chewed, held in mouth)

Use Pattern

People may use khaini from 3 to 30 times a day. A regular khaini user may carry a double-ended metal container, one side of which is filled with sun-dried tobacco and the other slightly moistened slaked lime.^{1,5}

Main Ingredients

Tobacco leaves, slaked lime (calcium hydroxide), and sometimes areca nut^{2,6}

Processing/Manufacturing

Custom-made, cottage industry, and commercial:

Khaini is usually prepared by the individual user from basic ingredients at the time of use, but commercially manufactured khaini is also available. To prepare khaini, users will use their thumbs to vigorously mix a small amount of dried tobacco leaves and slaked lime paste in the palm of the hand. Areca nut is sometimes added.² Additional information on the commercial manufacturing of khaini could not be located.

Khaini

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNA ^s *
		mg/g wet wt					
Khaini, † India	9.65–9.79	2.53–4.79	2.48–4.68	288–502	16,800–17,500	1,350–1,400	21,600–23,500

*Total TSNA^s represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).

†All khaini products were commercially manufactured.

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNA^s = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Stanfill et al. 2011 (6).

References

1. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf
2. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
3. International Institute for Population Sciences, Ministry of Health and Family Welfare, Government of India. Global Adult Tobacco Survey, India, 2009–2010. New Delhi: Ministry of Health and Family Welfare; Mumbai: International Institute for Population Sciences; 2010 [cited 21 June 2012]. Available from: http://whoindia.org/EN/Section20/Section25_1861.htm
4. Ministry of Health and Family Welfare (Bangladesh). Global Adult Tobacco Survey: Bangladesh report 2009. Dhaka, Bangladesh: World Health Organization, Country Office for Bangladesh; 2009. Available from: http://www.who.int/tobacco/surveillance/global_adult_tobacco_survey_bangladesh_report_2009.pdf
5. Public Health Foundation of India. Smokeless tobacco products commonly used in India [factsheet]. New Delhi: Public Health Foundation of India, Centre for Tobacco Control and Health Promotion. Available from: http://ctchp.org/index.php?option=com_content&view=article&id=171:smokeless-tobacco-products-commonly-used-in-india&catid=216&Itemid=159
6. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. *Tob Control*. 2011 May;20(3):e2. Epub 2010 Nov 25.

For additional information on khaini, please refer to the following chapters: Chapter 10: Smokeless Tobacco Use in the European Region, and Chapter 13: Smokeless Tobacco Use in the South-East Asia Region.

Kiwam

Kiwam (also spelled qiwam, qimam, khiwam, kimam) is a tobacco paste that is made from boiled and flavored tobacco leaves. The paste may be formed into granules or pellets. Kiwam is frequently used as the tobacco ingredient in betel quid (paan).^{1,2,3,4}



Photo courtesy of Clifford Watson, Centers for Disease Control and Prevention

Common Names

None

Brand Names

Avon, Kashmiri, Nauratan, Raj Ratan, Pradip

Main Geographic Locations

(WHO Region: Country)

South-East Asia Region: India, Bangladesh, Nepal^{1,2};

Eastern Mediterranean Region: Pakistan³

Prevalence and Demographics

No data are available on the prevalence of kiwam use. In 2004, it was reported that kiwam is used among upper socioeconomic groups in India and Bangladesh.¹

Mode of Absorption

Oral (chewed, held in mouth, chewed in betel quid)

Use Pattern

Kiwam may be used alone or inserted into a betel quid and chewed.² Additional information on patterns of kiwam use could not be located.

Main Ingredients

Tobacco, spices (cardamom, saffron, and/or aniseed), additives such as musk^{1,2}

Processing/Manufacturing

Commercial: Kiwam is commercially manufactured. The stalks, stems, and veins of tobacco leaves are removed, and then the remaining leaves are boiled, soaked in water, and flavored with powdered spices and other additives.¹ Once the tobacco has softened and broken down, the resulting pulp is mashed, strained, and dried into a paste.² Kiwam may be sold as a paste or in granule or pellet form.^{1,3}

Kiwam

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNA ^s *
		mg/g wet wt			ng/g wet wt		
Kiwam,† India	—	—	—	100–1,030	2,500–8,950	160–1,860	5,430–22,200

*Total TSNA^s represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).

†All kiwam products are commercially manufactured.

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNA^s = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Tricker and Preussmann 1989 (5) (more recent data was not available).

References

1. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf
2. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
3. Gupta PC, Ray CS. Smokeless tobacco and health in India and South Asia. *Respirology*. 2003;8(4):419–31.
4. Reddy KS, Gupta PC, editors. Report on tobacco control in India. New Delhi: Ministry of Health and Family Welfare, Government of India; 2004. Available from: http://www.who.int/fctc/reporting/Annex6_Report_on_Tobacco_Control_in_India_2004.pdf
5. Tricker AR, Preussmann R. The occurrence of *N*-nitroso compounds in kiwam tobacco. *Cancer Lett*. 1989;46(3):221–4.

Loose Leaf Chew

Three types of chewing tobacco are used in North America: loose leaf, plug, and twist. Although loose leaf is the most common form in the United States, the use of chewing tobacco has declined over the past few decades and is now uncommon.^{1,2}



Photo courtesy of Clifford Watson, Centers for Disease Control and Prevention

Common Names

Chew, chaw, chewing tobacco, spit tobacco

Brand Names

Red Man, Granger, J.D.'s Blend (*Swedish Match North America*); Levi Garrett, Morgan's, Taylor's Pride (*American Snuff Company*); Beech-Nut, Our Pride, Stoker (*National Tobacco Company*); Mail Pouch, Chattanooga Chew, Lancaster (*Swisher International*)

Main Geographic Location (WHO Region: Country)

Region of the Americas: United States¹

Prevalence and Demographics

Product-specific prevalence rates are not available for any countries where loose leaf chew is used. In the United States, chewing tobaccos are primarily used by men and are more common in rural areas and the South and Midwest.² In 2012, the U.S. prevalence of past-month smokeless tobacco use (including loose leaf tobacco, snuff, and other smokeless products) for those aged 12 years and older was 3.6% (7.1% of males and 0.4% of females).³ Although there are no statistics specifically on the prevalence of loose leaf tobacco use, sales of loose leaf chew represent 19.7% of smokeless tobacco sales in the United States.⁴

Mode of Absorption

Oral (chewed, sucked, held in mouth)

Use Patterns

A piece of tobacco 0.75 to 1 inch in diameter is either chewed or held in place. Saliva is usually spit out, but it can also be swallowed.¹

Main Ingredients

Leaf tobacco, sugar, and/or licorice^{1,5}

Processing/Manufacturing

Commercial: Loose leaf chew is commercially manufactured and usually sold in pouches. To manufacture loose leaf chew, loose cigar tobacco leaves are air-cured, stemmed, and cut or granulated to form small strips of shredded tobacco. Most brands are sweetened and flavored with sugar and licorice, accounting for loose leaf tobacco's high average sugar content (approximately 35% of its weight is sugar).^{1,5}

Loose Leaf Chew

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type*	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNAst
		mg/g wet wt		ng/g wet wt			
Loose Leaf (6), United States	5.64–5.98	4.87–7.04	0.03–0.06	238–306	942–2,830	20–90	1,550–4,100
Loose Leaf (1), United States	5.64–6.76	3.41–8.99	0.02–0.47	—	—	—	—

*All loose leaf products were commercially manufactured.

†Total TSNAst represent the sum of NNK, NNN, and NNAL (shown), *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAst = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Sources: Lawler et al. 2013 (6); International Agency for Research on Cancer 2007, table 5 (1).

References

1. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
2. Rodu B, Cole P. Smokeless tobacco use among men in the United States, 2000 and 2005. *J Oral Pathol Med.* 2009;38(7):545–50.
3. Substance Abuse and Mental Health Services Administration. Results from the 2012 National Survey on Drug Use and Health: detailed tables. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2013. Available from: <http://www.samhsa.gov/data/NSDUH/2012SummNatFindDefTables/DefTabs/NSDUH-DefTabsLOTsect2pe2012.htm#TopOfPage>
4. Maxwell J. The Maxwell report: the smokeless tobacco industry in 2010. Richmond, VA: John C. Maxwell, Jr.; 2011.
5. Hoffmann D, Djordjevic MV. Chemical composition and carcinogenicity of smokeless tobacco. *Adv Dent Res.* 1997;11(3):322–9.
6. Lawler TS, Stanfill SB, Zhang L, Ashley DL, Watson CH. Chemical characterization of domestic oral tobacco products: total nicotine, pH, unprotonated nicotine and tobacco-specific *N*-nitrosamines. *Food Chem Toxicol.* 2013;57:380–6. Epub 2013 Mar 19. doi 10.1016/j.fct.2013.03.011

For additional information on loose leaf chew, please refer to Chapter 9: Smokeless Tobacco Use in the Region of the Americas.

Mainpuri

Mainpuri is a mixture of tobacco, finely chopped areca nut, slaked lime, cloves, and camphor.^{1,2,3} It has a short shelf life and is mainly sold in the northern part of India in the Mainpuri district and nearby areas.¹ Mainpuri was also sold in neighboring Pakistan, although it was banned in 2011.



*Photo courtesy of Clifford Watson,
Centers for Disease Control and Prevention*

Common Names

Kapoori

Brand Names

None

Main Geographic Location

(WHO Region: Country)

South-East Asia Region: India (Uttar Pradesh)²

Prevalence and Demographics

Mainpuri is a very popular tobacco preparation in the Mainpuri district and surrounding areas of Uttar Pradesh.⁵ Recent data on the prevalence of mainpuri tobacco use in Uttar Pradesh or in India nationally are not available.

Mode of Absorption

Oral (chewed, chewed in betel quid, held in mouth)

Use Patterns

No information is available on use patterns.

Main Ingredients

Tobacco, slaked lime (calcium hydroxide), areca nut, camphor, cloves^{1,2}

Processing/Manufacturing

Cottage industry and custom-made: Mainpuri is manufactured by small cottage industries and prepared by individual vendors for sale. It is prepared by thoroughly mixing together tobacco, slaked lime, finely cut areca nut, and powdered camphor and cloves.³ Additional information on the preparation of mainpuri could not be located.

Mainpuri

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine mg/g wet wt	Free Nicotine mg/g wet wt	NNK	NNN	NNAL ng/g wet wt	Total TSNA ^s *
Mainpuri, Pakistan	7.65	1.28	0.38	6.05	106	25.9	219

*Total TSNA^s represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).
Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosornicotine;
NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNA^s = tobacco-specific nitrosamines; mg/g = milligram per gram;
ng/g = nanogram per gram.
Source: Stanfill et al. 2011 (4).

References

1. Chaudhry K. Is pan masala-containing tobacco carcinogenic? Natl Med J India. 1999;12(1):21-7.
2. Reddy KS, Gupta PC, editors. Report on tobacco control in India. New Delhi: Ministry of Health and Family Welfare, Government of India; 2004. Available from: http://www.who.int/fctc/reporting/Annex6_Report_on_Tobacco_Control_in_India_2004.pdf
3. Wahi P. The epidemiology of oral and oropharyngeal cancer: a report of the study in Mainpuri District, Uttar Pradesh, India. Bull World Health Organ. 1968;38(4):495-521.
4. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. Tob Control; 2011 May;20(3):e2. Epub 2010 Nov 25.
5. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf

Mawa

Mawa is a mixture of finely chopped ingredients consisting of areca nut shavings, slaked lime, and crushed sun-cured tobacco.^{1,2} It is about 95% areca nut by weight and is used in South Asia.^{1,3}



*Photo courtesy of Clifford Watson,
Centers for Disease Control and Prevention*

Common Names

None

Brand Names

None

Main Geographic Location

(WHO Region: Country)
South-East Asia Region: India¹

Prevalence and Demographics

Mawa is popular among men and young people, particularly in Gujarat, India.^{1,3} Between the 1970s and the 1990s, mawa chewing increased considerably in the Bhavnagar district and adjoining areas in Gujarat, particularly among young people. One survey in 1998 in this district found that 18.9% of men and 0.1% of women used mawa.⁴ A study conducted in 2007–2008 in urban Jamnagar, Gujarat, found that approximately 21% of people aged 13 and older chewed mawa.⁵ National data on the prevalence of mawa use in India are not available.

Mode of Absorption

Oral (chewed)

Use Pattern

Mawa is chewed for about 10 to 20 minutes¹ and may be used as many as 5 to 25 times per day.³

Main Ingredients

Tobacco, slaked lime, and areca nut¹

Processing/Manufacturing

Cottage industry and custom-made: Mawa is prepared by individual vendors for sale or is homemade by individual users. The process of making mawa consists of sprinkling small pieces of sun-cured areca nut with slaked lime and adding tobacco flakes. The mixture is rubbed together to combine the ingredients. Mawa is about 95% areca nut by weight and is sold in cellophane wrappers.³

Mawa

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine mg/g wet wt	Free Nicotine mg/g wet wt	NNK	NNN	NNAL ng/g wet wt	Total TSNAs*
Mawa, † Pakistan	8.31	0.16	0.11	4.47	65.5	3.98	96

*Total TSNAs represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).

†Data is given for custom-made mawa.

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAs = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Stanfill et al. 2011 (6).

References

1. International Agency for Research on Cancer. Betel-quid and areca-nut chewing and some areca-nut-derived nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 85. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2004. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol85/mono85.pdf>
2. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
3. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf
4. Gupta P, Sinor PN, Bhonsle RB, Pawar VS, Mehta HC. Oral submucous fibrosis in India: a new epidemic? Natl Med J India. 1998;11(3):113–6.
5. Joshi U, Modi B, Yadav S. A study on prevalence of chewing form of tobacco and existing quitting patterns in urban population of Jamnagar, Gujarat. Indian J Community Med. 2010;35(1):105–8.
6. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. Tob Control. 2011 May;20(3):e2. Epub 2010 Nov 25.

For additional information on mawa, please refer to Chapter 13: Smokeless Tobacco Use in the South-East Asia Region.

Mishri

Mishri (also known as masherior or misherri) is a dry, powdered tobacco product that is usually homemade or prepared by a vendor. Mishri is commonly believed to clean teeth.¹



Photo courtesy of World Health Organization South-East Asia Regional Office and Dharendra N. Sinha

Common Names

None

Brand Names

None

Main Geographic Location (WHO Region: Country)

South-East Asia Region: India^{2,3}

Prevalence and Demographics

Mishri is predominantly used by women and is more common among lower socioeconomic groups.⁴ Many users begin using mishri as children. While there are no national data or recent prevalence estimates, various studies of different regions from the 1970s to the present have estimated that mishri use in India has ranged between 17% and 44% of women and between 1% and 23% of men.^{3,5,6,7} In 2009–2010, 5% of all individuals aged 15 years and over (3.3% of males and 6.3% of females) in India consumed oral tobacco including mishri, snuff, gul, or gudakhu.⁸

Mode of Absorption

Oral (sucked, applied to teeth and gums, teeth cleaning)¹

Use Pattern

Mishri is rubbed on the teeth and gums, often for the purpose of cleaning the teeth. It is generally used twice a day, but users who become addicted may apply it more frequently or hold it in their mouths.^{1,3,4}

Main Ingredients

Tobacco¹

Processing/Manufacturing

Custom-made and cottage industry: Mishri is usually prepared at home by individual users, but it can also be prepared by vendors for sale and bought in the market under various names.⁵ No other ingredients besides tobacco are used in the preparation of mishri. Tobacco is baked on a hot metal plate until it is toasted or partially burnt and is uniformly black. It is then powdered.¹

Mishri

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine (9)	Free Nicotine (9)	NNK (10)	NNN (10)	NNAL	Total TSNAs
		mg/g wet wt				ng/g wet wt	
Mishri,* India	6.54	2.73	0.09	4,210	870	—	—

*Chemical measurements from both sources are for Shahin mishri.

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = N'-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAs = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Sources: Gupta and Sreevidya 2004 (9); Stepanov et al. 2005 (10).

References

1. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific N-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
2. Rani M, Bonu S, Jha P, Nguyen SN, Jamjoum L. Tobacco use in India: prevalence and predictors of smoking and chewing in a national cross sectional household survey. *Tob Control*. 2003;12(4):e4.
3. Reddy KS, Gupta PC, editors. Report on tobacco control in India. New Delhi: Ministry of Health and Family Welfare, Government of India; 2004. Available from: http://www.who.int/fctc/reporting/Annex6_Report_on_Tobacco_Control_in_India_2004.pdf
4. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf
5. Pratinidhi A, Gandham S, Shrotri A, Patil A, Pardeshi S. Use of "mishri": a smokeless form of tobacco during pregnancy and its perinatal outcome. *Indian J Community Med*. 2010;35(1):14-8.
6. Gupta PC. Survey of sociodemographic characteristics of tobacco use among 99,598 individuals in Bombay, India using handheld computers. *Tob Control*. 1996;5(2):114-20.
7. Mehta FS, Gupta PC, Daftary DK, Pindborg JJ, Choksi SK. An epidemiologic study of oral cancer and precancerous conditions among 101,761 villagers in Maharashtra, India. *Int J Cancer*. 1972;10(1):134-41.
8. International Institute for Population Sciences, Ministry of Health and Family Welfare, Government of India. Global Adult Tobacco Survey, India, 2009-2010. New Delhi: Ministry of Health and Family Welfare; Mumbai: International Institute for Population Sciences; 2010 [cited 21 June 2012]. Available from: http://whoindia.org/EN/Section20/Section25_1861.htm
9. Gupta PC, Sreevidya S. Report on smokeless tobacco products in India. Mumbai: Action Council Against Tobacco; 2004.
10. Stepanov I, Hecht SS, Ramakrishnan S, Gupta PC. Tobacco-specific nitrosamines in smokeless tobacco products marketed in India. *Int J Cancer*. 2005;116(1):16-9

For additional information on mishri, please refer to Chapter 13: Smokeless Tobacco Use in the South-East Asia Region.

Moist Snuff

(Dip)

Moist snuff is a damp, finely ground tobacco product that is more commonly used in Western countries than elsewhere. Moist snuff, also referred to as dip, is the most common form of smokeless tobacco in the United States and Canada.^{1,2} This factsheet only describes commercial moist snuff. Other types of moist snuff are used in many countries around the world.



Common Names

Dip, spit tobacco (sometimes called “chew,” even though it is not chewing tobacco)

Brand Names

Copenhagen, Skoal, Red Seal, Husky (*U.S. Smokeless Tobacco Company*), Grizzly, Kodiak (*American Snuff Company*), Kayak, Redwood, Gold River, Silver Creek, Cooper, Silverado (*Swisher International*), Red Man, Timber Wolf, Longhorn (*Swedish Match*)

Main Geographic Locations (WHO Region: Country)

Region of the Americas: United States, Canada,^{1,2} Mexico³; **African Region:** South Africa^{1,4}

Prevalence and Demographics

Moist snuff is the most commonly used form of smokeless tobacco in the United States, Canada, and Mexico.^{1,2,3} In the United States, it is primarily used by young adult white men living in the South.⁵ Although no statistics specifically on the prevalence of moist snuff use are available, sales of moist snuff represented 78.2% of U.S. smokeless tobacco sales in 2010.⁶ The prevalence of smokeless tobacco use in the United States in 2012 was 3.6% among people age 12 and older (7.1% of males and 0.4% of females).⁷ In Canada, smokeless tobacco use is low—1% of adult men use any type of smokeless tobacco⁸—and sales of moist snuff make up more than 80% of all smokeless tobacco sales.²

Mode of Absorption

Oral (held in mouth, sucked)

Use Pattern

Moist snuff is commonly used loose, but can also come in small, ready-to-use pouches. It is usually held in the mouth for about 30 minutes. Saliva is usually spit out, but it can be swallowed.¹

Main Ingredients

Tobacco, flavorings, inorganic salts, humectants¹

Processing/Manufacturing

Commercial: Moist snuff is commercially manufactured. Tobacco leaf, stems, and seeds are air- and/or fire-cured, then processed into fine particles (“fine cut”) or strips (“long cut”).¹ This process ferments the tobacco, which produces more cancer-causing nitrosamines than snus manufacturing⁹ (snus is often considered a different type of moist snuff). Moist snuff has a moisture content of roughly 20–60% by weight and is available in a variety of flavors, such as mint and fruit flavors. The tobacco is usually sold loose, but can also be packaged in small, tea bag-like pouches called sachets.¹

Moist Snuff (Dip)

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type*	pH	Total Nicotine mg/g wet wt	Free Nicotine mg/g wet wt	NNK	NNN	NNAL ng/g wet wt	Total TSNAst
Moist Snuff (10), United States	5.54–8.62	4.42–25.03	0.01–7.81	382–9,950	2,204–42,554	21–1,412	4,874–90,024
Moist Snuff (1), United States	5.49–8.38	7.06–24.29	0.03–8.57	—	—	—	—

*Moist snuff products were commercially manufactured.

†Total TSNAst represent the sum of NNK, NNN, and NNAL (shown), *N*'-nitrosoanatabine and *N*'-nitrosoanabasine (not shown).

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N*'-nitrososnicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAst = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Sources: Richter et al. 2008 (10); International Agency for Research on Cancer 2007, table 7 (1).

References

- International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
- Euromonitor International. Country report: smokeless tobacco in Canada. August 2012. Available by subscription from <http://www.euromonitor.com>.
- Euromonitor International. Country report: smokeless tobacco in Mexico. October 2012. Available by subscription from <http://www.euromonitor.com>.
- Ayo-Yusuf OA, Swart TJ, Pickworth WB. Nicotine delivery capabilities of smokeless tobacco products and implications for control of tobacco dependence in South Africa. *Tob Control*. 2004;13(2):186–9.
- Rodu B, Cole P. Smokeless tobacco use among men in the United States, 2000 and 2005. *J Oral Pathol Med*. 2009;38(7):545–50.
- Maxwell J. The Maxwell report: the smokeless tobacco industry in 2010. Richmond, VA: John C. Maxwell, Jr.; 2011.
- Substance Abuse and Mental Health Services Administration. Results from the 2012 National Survey on Drug Use and Health: detailed tables. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2013. Available from: <http://www.samhsa.gov/data/NSDUH/2012SummNatFindDetTables/DefTabs/NSDUH-DefTabsLOTsect2pe2012.htm#TopOfPage>
- Health Canada. Canadian tobacco use monitoring survey, 2010. User guide. Ottawa, Ontario, Canada: Health Canada; 2010.
- Hatsukami DK, Ebbert JO, Feuer RM, Stepanov I, Hecht SS. Changing smokeless tobacco products: new tobacco delivery systems. *Am J Prev Med*. 2007;33(6Suppl):S368–78.
- Richter P, Hodge K, Stanfill S, Zhang L, Watson C. Surveillance of moist snuff: total nicotine, moisture, pH, un-ionized nicotine, and tobacco-specific nitrosamines. *Nic Tob Res*. 2008 Nov;10(11):1645–52.

For additional information on moist snuff, please refer to the following chapters: Chapter 9: Smokeless Tobacco Use in the Region of the Americas, and Chapter 12: Smokeless Tobacco Use in the African Region.

Nass

Nass (also known as naswar) is a multinational product made of locally grown tobacco, oil, an alkaline modifier such as ash or slaked lime, and other ingredients according to regional preference.^{1,2} Nass/nasway is one of the most common types of smokeless tobacco used in Pakistan, the United Arab Emirates, Uzbekistan, and Kyrgyzstan.^{1,3} In Uzbekistan and Kyrgyzstan, a product known as nasway contains the same main ingredients as nass. Although there is insufficient published information to determine if nass and nasway are the same product, for the purposes of this factsheet, they are assumed to be essentially the same.




Photo courtesy of Clifford Watson, Centers for Disease Control and Prevention

Common Names

Naswar (Pakistan); niswar (United Arab Emirates); nass (Iran); nasway, nasvay (Kyrgyzstan, Uzbekistan)

Brand Names

None

Main Geographic Locations

(WHO Region: Country)

Eastern Mediterranean Region: Pakistan, Iran, Afghanistan, United Arab Emirates^{1,4}; **African Region:** South Africa^{1,4}; **European Region:** Turkmenistan, Kyrgyzstan, Uzbekistan^{5,6}

Prevalence and Demographics

In Kyrgyzstan, according to a 2011 World Health Organization report, 3.4% of all adults (7.0% of men and 0.3% of women) use nasvay.⁵ In Uzbekistan, in a 2002 survey of adults between the ages of 15 and 60, only 0.4% of women reported ever using nasway in their lifetime, compared to 37.9% of men. Thus, among men in Uzbekistan, using nasway is as common as cigarette smoking.⁷ Studies in Turkmenistan in 1993 reported that 12% of adults used nass.⁶

Mode of Absorption

Oral (chewed, sucked, held in mouth)

Use Pattern

Nass is usually rolled into a ball and placed under the tongue or in the cheek.^{1,4,8} It is held in the mouth and sucked for 10 to 15 minutes or may be chewed slowly.²

Main Ingredients

Nass: tobacco, ash, cotton or sesame oil, water, and sometimes lime or gum^{2,9}

Naswar, niswar, nasway: tobacco, slaked lime (calcium hydroxide), ash, oil or butter, indigo or other coloring agent, water, and sometimes flavorings such as cardamom and menthol^{1,2,4,7,8,9}

Processing/Manufacturing

Cottage industry and custom-made: Nass may be made domestically by individual users or prepared by local, small-scale cottage manufacturers.^{4,9} Sun- and heat-dried tobacco leaves (often *N. rustica*), slaked lime, ash from tree bark, flavorings (e.g., cardamom and menthol), and coloring agents (e.g., indigo) are mixed together with a heavy wooden mallet. Water is added and the mixture is usually rolled into balls.^{1,4} The product is then packed into small polyethylene bags for sale.⁴

Nass

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine mg/g wet wt	Free Nicotine mg/g wet wt	NNK	NNN	NNAL	Total TSNA ^s *
Naswar, [†] Pakistan	8.76–9.14	10.5–14.2	8.84–13.2	29.4–309	363–545	8.6–104	478–1,380
Nasway, [†] Uzbekistan	8.43	8.89	6.36	88.3	628	10.5	1,100

*Total TSNA^s represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).

[†]Naswar/nasway products were produced by cottage industry.

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosonornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNA^s = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Stanfill et al. 2011 (10).

References

1. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
2. Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR). Health effects of smokeless tobacco products. Brussels: European Commission; 2008 [cited 2012 Oct 25]. Available from: http://ec.europa.eu/health/archive/ph_risk/committees/04_scenihr/docs/scenihr_o_013.pdf
3. The National. UAE: police close chewing tobacco factories to clean up towns and cities. The National. 2009 March 31. [cited 2012 Aug 27]. Available from: http://www.daijworld.com/news/news_disp.asp?n_id=58384&n_tit=UAE%3A+Police+Close+Chewing+Tobacco+Factories+to+Clean+Up+Towns+%26+Cities
4. Zakiullah, Saeed M, Muhammad N, Khan SA, Gul F, Khuda F, et al. Assessment of potential toxicity of a smokeless tobacco product (naswar) available on the Pakistani market. *Tob Control*. 2012; 21(4):396–401.
5. World Health Organization. WHO report on the global tobacco epidemic, 2011: warning about the dangers of tobacco. Geneva: World Health Organization; 2011. Available from: http://www.who.int/tobacco/global_report/2011/en/
6. World Health Organization. Tobacco or health: a global status report. Geneva: World Health Organization; 1997.
7. Ministry of Health (Uzbekistan). Uzbekistan Health Examination Survey, 2002. Calverton, MD: Uzbekistan Ministry of Health, Analytical and Information Center; Uzbekistan Ministry of Macroeconomics and Statistics, State Department of Statistics; and ORC Macro; 2004. Available from: http://pdf.usaid.gov/pdf_docs/PNACX976.pdf
8. Hammond SK. Global patterns of nicotine and tobacco consumption. In: Henningfield JE, London ED, Pogun S, editors. Nicotine psychopharmacology. Handbook of experimental pharmacology. Vol. 192. Berlin: Springer; 2009. p. 3–28.
9. Gupta PC, Ray CS. Smokeless tobacco and health in India and South Asia. *Respirology*. 2003;8(4):419–31.
10. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. *Tob Control*. 2011 May;20(3):e2. Epub 2010 Nov 25.

For additional information on nass/nasway, please refer to the following chapters: Chapter 10: Smokeless Tobacco Use in the European Region, and Chapter 11: Smokeless Tobacco Use in the Eastern Mediterranean Region.

Plug

Plug, one of three types of chewing tobacco sold in North America, is considered the oldest form of chewing tobacco. Plug tobacco is sweetened with sugar and licorice and pressed into a brick shape. It is sold as either “moist” or “firm” depending on the amount of moisture in the product.¹



*Photo courtesy of Clifford Watson,
Centers for Disease Control and Prevention*

Common Names

Chew, chaw, chewing tobacco, spit tobacco

Brand Names

Red Man, Days Work, Apple, Brown, Natural Leaf, Union Standard, Tinsley, WNT (*Swedish Match North America*), Levi Garrett, Taylors Pride, Cannon Ball (*American Snuff Company*)

Main Geographic Location (WHO Region: Country)

Region of the Americas: United States¹

Prevalence and Demographics

Chewing tobaccos are primarily used by men and are more commonly used in rural areas and in the Southern and Midwestern United States.² Plug use has declined over the past century and is rare today in most regions of the United States.¹ Although there are no recent statistics specifically on the prevalence of plug use, in 2010 sales of plug represented only 0.5% of all smokeless tobacco sales in the United States.³

Mode of Absorption

Oral (chewed, sucked, held in mouth)

Use Pattern

A piece of plug is cut off and chewed or held between the cheek and gum. Saliva is usually spit out, but it can also be swallowed.¹

Main Ingredients

Burley and bright tobacco or cigar tobacco leaves, licorice, and sugar¹

Processing/Manufacturing

Commercial: Plug is commercially manufactured. Heavier grades of tobacco leaves are picked from the top of the plant and stems are removed. The tobacco is immersed in a mixture of licorice and sugar, pressed into a plug, wrapped in fine tobacco leaves, and pressed into bricks or flat blocks. Moist plug tobacco has at least 15% moisture by weight, whereas firm plug has less than 15% moisture. Plug also has a high average sugar content (approximately 25% of its weight).¹

Plug

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type*	pH	Total Nicotine mg/g wet wt	Free Nicotine mg/g wet wt	NNK	NNN	NNAL	Total TSNAst
Plug (4), United States	5.10–5.95	5.12–15.1	0.01–0.04	340–941	2,920–5,140	11–188	4,090–7,750
Plug (1), United States	5.07–5.95	6.18–20.43	0.02–0.08	—	—	—	—

*All loose leaf products were commercially manufactured.

†Total TSNAst represent the sum of NNK, NNN, and NNAL (shown), *N*'-nitrosoanatabine and *N*'-nitrosoanabasine (not shown).

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N*'-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAst = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Sources: Lawler et al. 2013 (4); International Agency for Research on Cancer 2007, table 5 (1).

References

1. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
2. Rodu B, Cole P. Smokeless tobacco use among men in the United States, 2000 and 2005. *J Oral Pathol Med.* 2009;38(7):545–50.
3. Maxwell J. The Maxwell report: the smokeless tobacco industry in 2010. Richmond, VA: John C. Maxwell, Jr.; 2011.
4. Lawler TS, Stanfill SB, Zhang L, Ashley DL, Watson CH. Chemical characterization of domestic oral tobacco products: total nicotine, pH, unprotonated nicotine and tobacco-specific *N*-nitrosamines. *Food Chem Toxicol.* 2013;57:380–6. Epub 2013 Mar 19. doi 10.1016/j.fct.2013.03.011

Rapé

Rapé is a type of dry snuff used in Brazil. Aboriginal groups may mix it with ashes from particular trees and use it for medicinal purposes.¹ Rapé is usually inhaled through the nose.



Photo courtesy of Clifford Watson,
Centers for Disease Control and Prevention

Common Names

None

Commercial Names

Moeda, Caratinga, Guarany

Main Geographic Location

(WHO Region: Country)

Region of the Americas: Brazil¹

Prevalence and Demographics

Published information on rapé is limited, and data on the prevalence of rapé use are not available. The 2001 Brazilian Global Adult Tobacco Survey found that about 0.4% of adults aged 15 years and older use any type of smokeless tobacco.² Anecdotal evidence suggests that rapé is primarily used in rural areas and small towns, or by aboriginals in the Amazon rainforest, where its use has cultural significance.

Mode of Absorption

Nasal

Use Pattern

Rapé is inhaled through the nose. Some aboriginal groups prepare their own rapé. The Kaxinawás

Indians from Acre in Brazil prepare a half portion of rapé and half portion of ashes from wood, usually from the paricá tree (*Schizolobium amazonicum*). They consume rapé by having one person use a large “V”-shaped straw to blow the dust into their partner’s nose. They may use it for medicinal purposes.¹

Main Ingredients

Dried tobacco leaf, flavorings such as tonka bean (*Dipteryx odorata*), clove, cinnamon powder, and camphor, and in some cases, ashes from select trees

Processing/Manufacturing

Cottage industry and custom-made: Rapé is produced locally on small farms, in small tobacco industries, or by Indians. The dried tobacco leaf is ground and carefully toasted. Rapé is sometimes toasted with other ingredients such as spices, herbs, and ashes, and the toasted product is then sifted into a very fine dust.

Chemical Measurements

No data available

References

1. Lagrou EM. Xamanismo e representação entre os Kaxinawás. In: Langdon EJM, editor. Xamanismo no Brasil: novas perspectivas. Florianópolis: Editora da UFSC; 1996. p. 197–231. Portuguese.
2. Centers for Disease Control and Prevention. Global Adult Tobacco Survey (GATS), 2008–2010: percentage of adults who currently use smokeless tobacco. Global Tobacco Surveillance System data [Internet database]. Atlanta: Centers for Disease Control and Prevention [no date]. Available from: <http://nccd.cdc.gov/GTSSData/default/default.aspx>

For additional information on rapé, please refer to Chapter 9: Smokeless Tobacco Use in the Region of the Americas.

Red Toothpowder

Red toothpowder contains finely powdered tobacco, herbs, flavoring agents, and other ingredients.¹ It is red in color and is usually used in India to clean teeth.^{1,2} In 1992, India banned the use of tobacco as an ingredient in dental products.^{2,3} Brands of red toothpowder have stopped listing tobacco as an ingredient, but lab results show that some still contain tobacco and measureable amounts of nicotine.^{2,3,4,5}



Photo courtesy of World Health Organization South-East Asia Regional Office and Dharendra N. Sinha

Common Names

Lal dant manjan

Brand Names

Dabur, Baidhyanath

Main Geographic Location

(WHO Region: Country)

South-East Asia Region: India^{1,2}

Prevalence and Demographics

Red toothpowder is used by men and women of all ages, as well as children.² There are no national statistics on the prevalence of use of red toothpowder by adults. The 2004 Global Youth Tobacco Survey, which surveys school students aged 13–15 years, found that reported prevalence of red toothpowder use ranged from 2% to 49% across various regions in India.³

Mode of Absorption

Oral (teeth cleaning)

Use Patterns

Red toothpowder is used to clean teeth (as a dentifrice).^{1,2}

Main Ingredients

Fine red tobacco powder, herbs, flavorings.¹ Additional plant-related ingredients such as ginger, pepper, and camphor, among others, may be used.^{6,7}

Processing/Manufacturing

Commercial: Red toothpowder is commercially manufactured and is often marketed as an herbal dental care product.¹ Additional information on the manufacturing of red toothpowder is not available.

Red Toothpowder

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNAs*
		mg/g wet wt			ng/g wet wt		
Red Toothpowder,† India	5.75–6.71	4.47–5.09	0.03–0.21	—	—	—	—

*Study did not measure NNK, NNN, or NNAL TSNAs.

†All red toothpowder products were commercially manufactured.

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = N'-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAs = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Gupta and Sreevidya 2004 (5).

References

1. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific N-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
2. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf
3. Sinha DN, Gupta PC, Pednekar MS. Use of tobacco products as dentifrice among adolescents in India: questionnaire study. *Brit Med J*. 2004;328(7435):323–4.
4. Reddy KS, Gupta PC, editors. Report on tobacco control in India. New Delhi: Ministry of Health and Family Welfare, Government of India; 2004. Available from: http://www.who.int/fctc/reporting/Annex6_Report_on_Tobacco_Control_in_India_2004.pdf
5. Gupta PC, Sreevidya S. Report on smokeless tobacco products in India. Mumbai: Action Council Against Tobacco; 2004.
6. Shre Baidyanath Ayurved Bhawan PVT. Featured products – dant manjan lal [Internet]. Kolkata: Baidyanath; c2009 [cited 2012 Oct 3]. Available from: http://www.baidyanath.com/our_products.php?fea=5
7. Dabur India. Dabur products: Dabur lal dant manjan contains [Internet]. Ghaziabad, India; Dabur; [no date]. [updated 2012 Oct 3; cited 2012 Oct 3]. Available from: <http://www.dabur.com/Products-Health%20Care-Red%20Tooth%20Powder>

Shammah

Shammah is a multinational smokeless tobacco product in powder or paste form, which can be either greenish-yellow (“white” shammah) or brownish-black (“black” shammah). It is made from powdered tobacco, slaked lime, ash, and black pepper and mainly used in Yemen, Saudi Arabia, and Algeria.¹



Photo courtesy of Clifford Watson, Centers for Disease Control and Prevention

Common Names

El-Shama, bajeli, haradi, sharaci, black shammah (Yemen); al-shammah (Saudi Arabia); chemma (Algeria)

Brand Names

None

Main Geographic Locations

(WHO Region: Country)

Eastern Mediterranean Region: Saudi Arabia, Yemen^{1,2,3}; **African Region:** Algeria^{1,4}

Prevalence and Demographics

In Yemen in 2003, 10.7% of the population aged 10 years and older used shammah (15.1% of males and 6.2% of females).² Shammah was used by males and females of all ages, including adolescents. Shammah use increases with age and is more common in rural than urban areas.² In Algeria, shammah (or chemma) is commonly consumed by men across all social groups.⁴ Information on prevalence of shammah use in Algeria and Saudi Arabia is not available.

Mode of Absorption

Oral (sucked, held in mouth)

Use Pattern

Shammah is placed between the gum and lower lip or cheek.^{1,2} In Algeria, users may wrap shammah in paper before putting it in the mouth.⁴

Main Ingredients

Tobacco, slaked lime, ash, black pepper, oil, flavorings, bombosa (sodium carbonate)^{1,2}

Processing/Manufacturing

Cottage industry and custom-made: Shammah is usually prepared by small cottage industries or by local individual vendors for sale. The tobacco leaves are sun dried, pulverized with bombosa (sodium carbonate), and combined with other ingredients such as slaked lime, ash, black pepper, oil, and other flavorings. Shammah can be sold as a wet or dry product. To prepare wet shammah, such as black shammah, a water solution of bombosa is used rather than the dry powder alone.

Shammah

Chemical Measurements

No data available

References

1. Scheifele C, Nassar A, Reichart PA. Prevalence of oral cancer and potentially malignant lesions among shammah users in Yemen. *Oral Oncol.* 2007;43(1):42–50.
2. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific N-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/index.php>
3. Ministry of Public Health and Population (Yemen). The 2003 Family Health Survey sample in Yemen. [no city]: Republic of Yemen, Ministry of Public Health and Population; [no date] [cited 2012 Oct 26]. Available from: http://www.mophp-ye.org/arabic/docs/Familyhealth_english.pdf
4. Euromonitor International. Country report: smokeless tobacco in Algeria. November 2011. Available by subscription from: <http://www.euromonitor.com>.

For additional information on shammah, please refer to the following chapters: Chapter 11: Smokeless Tobacco Use in the Eastern Mediterranean Region, and Chapter 12: Smokeless Tobacco Use in the African Region.

Snus

Snus is a traditional Swedish smokeless tobacco product. It is made from moist, finely ground tobacco and usually contains lower levels of tobacco-specific *N*-nitrosamines (TSNAs) than most oral tobacco products because it is pasteurized rather than fermented.^{1,2} Although the use of snus is spreading to other regions of the world, in 1992 the sale of snus was banned in all countries in the European Union (EU) except Sweden.²



Photo courtesy of Clifford Watson,
Centers for Disease Control and Prevention

Common Names

None

Brand Names

General, Catch, Ettan, Grovsnus, Göteborgs Rapé, Kronan (*Swedish Match*); Lucky Strike, Pall Mall, du Maurier (*British American Tobacco*); Camel (*R.J. Reynolds*); Marlboro (*Philip Morris*); Skoal (*U.S. Smokeless Tobacco Company*); Knox, Skruf (*Imperial Tobacco*), Tobaccorette

Main Geographic Locations (WHO Region: Country)

European Region: Sweden, Norway, Iceland, Finland, Denmark^{3,4}; **Region of the Americas:** United States, Canada, Brazil³; **African Region:** South Africa⁵

Prevalence and Demographics

In Sweden in 2009, the prevalence of daily snus use among adults aged 16 years and older was 19% for males and 4% for females.⁶ In Norway, which is not a member of the EU and therefore can legally sell snus, the prevalence of daily use in 2009 was 6% among adults aged 16 years and older (11% of males and 1% of females).⁴ In 2010, 5.1% of U.S. adults aged 18 years and older had ever tried snus (8.5% of males and 2% of females), yet less than 1% currently used snus.⁷ There is little information on prevalence of use in most of the other countries where snus is available. Although snus has been used in Europe's Nordic region for many years, it was only introduced relatively recently into the North American and South African markets, and therefore the prevalence of

snus use may increase in these regions as this product becomes more widely available.^{5,7}

Mode of Absorption

Oral (held in mouth)

Use Patterns

Snus can either be packaged into small, ready-to-use sachets or sold in loose tobacco form. One portion of snus is usually held in the mouth for 30 minutes or more and does not require chewing, sucking, or spitting. In Sweden the average user keeps snus in their mouth for 11 to 14 hours per day.³

Main Ingredients

Tobacco, moisturizers, sodium carbonate, salt (sodium chloride), sweeteners, flavorings^{3,8}

Processing/Manufacturing

Commercial: Snus is commercially manufactured, but processing and manufacturing vary across regions and countries. In Sweden, finely ground air-cured tobacco is mixed with salts, water, and flavoring. Snus is similar to snuff, but snuff is fermented, which can increase the formation of TSNAs. Snus goes through a heat treatment process that pasteurizes the tobacco to kill off bacteria that aid in the formation of TSNAs.² During pre-sale storage, snus is kept cold to keep it "fresh" and to prevent more nitrosamines from forming.⁹ However, variations in processing and manufacturing can produce variations in TSNAs.^{3,8,9} Snus is either sold loose or portion packed in small tea bag-like pouches, or sachets. Snus comes in a variety of flavors.³

Snus

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type*	pH	Total Nicotine mg/g wet wt	Free Nicotine mg/g wet wt	NNK	NNN	NNAL	Total TSNA [†] ng/g wet wt
RJ Reynolds (10), United States	7.55–7.70	8.97–11.3	2.51–3.69	84–146	369–425	20–21	761–884
Swedish Match (8), Sweden	6.61–7.21	7.76–15.2	0.29–2.03	84.5–105	267–345	8.57–13.1	601–723
British American Tobacco (8), South Africa	6.48–7.02	13.4–17.2	0.47–1.19	171–275	925–1,440	18.6–30.4	1,720–2,700
Tobacco-rette (8), South Africa	6.56	15.0	0.49	1,340	2,950	84.2	5,850

*All snus products were commercially manufactured; manufacturer (associated products): RJ Reynolds (Camel snus); Swedish Match (General snus; Catch Peppermint snus); and British American Tobacco (Peter Stuyvesant snus; Lucky Strike snus).
[†]Total TSNA represents the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).
 Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNA = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.
 Sources: Stanfill et al. 2011 (8); Lawler et al. 2013 (10).

References

1. Biener L, McCausland K, Curry L, Cullen J. Prevalence of trial of snus products among adult smokers. *Am J Public Health*. 2011;101(10):1874–6. Available from: <http://ajph.aphapublications.org/cgi/content/abstract/101/10/1874>
2. Fagerström KO, Schildt EB. Should the European Union lift the ban on snus? Evidence from the Swedish experience. *Addiction*. 2003;98(9):1191–5.
3. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
4. Helleve A, Weisæth AH, Lindbak RL, editors. *Tal om tobakk 1973–2009* [Tobacco use 1973–2009]. Oslo: Norwegian Directorate of Health; 2011. Available from: <http://www.helsedirektoratet.no/publikasjoner/tal-om-tobakk-1973-2009/Publikasjoner/tal-om-tobakk-1973-2009-nynorsk.pdf>. Norwegian.
5. Ayo-Yusuf OA, Swart TJP, Pickworth WB. Nicotine delivery capabilities of smokeless tobacco products and implications for control of tobacco dependence in South Africa. *Tob Control*. 2004;13(2):186–9.
6. Forey B, Hamling J, Hamling J, Lee P. International smoking statistics: a collection of worldwide historical data: Sweden. Sutton, UK: PN Lee Statistics & Computing; 2011 Oct. Available from: http://www.pnlee.co.uk/Downloads/ISS/ISS-Sweden_111024.pdf
7. McMillen R, Maduka J, Winickoff JP. Use of emerging tobacco products in the United States. *J Environ Public Health*. 2012; Article ID 989474. Epub 2012 May 10.
8. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. *Tob Control*. 2011 May;20(3):e2. Epub 2010 Nov 25.
9. Hatsukami DK, Ebbert JO, Feuer RM, Stepanov I, Hecht SS. Changing smokeless tobacco products: new tobacco delivery systems. *Am J Prev Med*. 2007;33(6S):S368–78.
10. Lawler TS, Stanfill SB, Zhang L, Ashley DL, Watson CH. Chemical characterization of domestic oral tobacco products: total nicotine, pH, unprotonated nicotine and tobacco-specific *N*-nitrosamines. *Food Chem Toxicol*. 2013;57:380–6. Epub 2013 Mar 19. doi 10.1016/j.fct.2013.03.011

For additional information on snus, please refer to the following chapters: Chapter 9: *Smokeless Tobacco Use in the Region of the Americas*; Chapter 10: *Smokeless Tobacco Use in the European Region*; and Chapter 12: *Smokeless Tobacco Use in the African Region*.

Tapkeer

Tapkeer (tapkir) is a form of dry powdered snuff that may be used for teeth cleaning. It is similar to other teeth-cleaning products such as mishri.¹

Common Names

Bajjar (Gujarat, India)

Brand Names

None

Main Geographic Location

(WHO Region: Country)

South-East Asia Region: India¹

Prevalence and Demographics

In Goa, Maharashtra, Gujarat, and eastern India, tapkeer is widely used by people of lower socioeconomic status and is more commonly used by women than men.^{1,2} Although there are no national or recent statistics specifically on tapkeer use, historical data indicate that 14% of women and 1% of men use tapkeer in Gujarat, India.³ In 2009–2010, 4.7% of adults aged 15 and older (3.3% of males and 6.3% of females) reported using at least one type of applied oral tobacco product, including tapkeer, mishri, or gudakhu.⁴

Mode of Absorption

Oral (teeth cleaning, held in mouth), Nasal

Use Pattern

In India, tapkeer is rubbed on the teeth and gums to clean teeth. Because it is addictive, users tend to use it several times a day.^{1,2}

Main Ingredients

Tobacco¹

Processing/Manufacturing

Custom-made: In India, tapkeer is frequently prepared by individual users at home by roasting and then powdering the tobacco.¹

Chemical Measurements

No data available

References

1. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/index.php>
2. Gupta PC, Ray CS. Smokeless tobacco and health in India and South Asia. *Respirology*. 2003;8(4):419–31.
3. Bhonsle RB, Murli PR, Gupta PC. Tobacco habits in India. In: Gupta PC, Hamner JE III, Murli PR, editors. Control of tobacco-related cancers and other diseases. Proceedings of an international symposium, TIFR. Bombay, January 15–19, 1990. Bombay: Oxford University Press; 1992. p. 25–46.
4. International Institute for Population Sciences, Ministry of Health and Family Welfare, Government of India. Global Adult Tobacco Survey, India, 2009–2010. New Delhi: Ministry of Health and Family Welfare; Mumbai: International Institute for Population Sciences; 2010 [cited 21 June 2012]. Available from: http://whoindia.org/EN/Section20/Section25_1861.htm

Tobacco Leaf

Raw dried tobacco leaf may be chewed alone, but it is frequently used in betel quid in India, Bangladesh, and Myanmar.¹



Common Names

Sada pata; chadha (Assam, India)

Brand Names

None

Main Geographic Locations (WHO Region: Country)

South-East Asia Region: India, Bangladesh, Myanmar, Bhutan^{1,2}

Prevalence and Demographics

In Bangladesh in 2009, 1.8% of adults aged 15 and older chewed sada pata (plain tobacco flakes) alone (2% of males, 1.6% of females); this figure does not include adults who used tobacco leaf in betel quid.³ National prevalence data on the use of tobacco leaf are not available for other countries. In Bangladesh and Myanmar, betel quid chewers of low socioeconomic status often put tobacco leaf in their quid.¹

Mode of Absorption

Oral (chewed, chewed in betel quid or other custom-made product)

Use Pattern

In India, a regular user chews about a 15-cm piece of the tobacco leaf per day.¹

Main Ingredients

Tobacco leaf¹

Processing/Manufacturing

Custom-made: Raw tobacco leaf is usually dried and left unprocessed. It can be powdered, flaked, or sold in bundles of several long strands (about 115 cm long and 5 cm thick). A regular user consumes one 15-cm piece of the strand per day.¹

Tobacco Leaf

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNA*s*
		mg/g wet wt					
Tobacco Leaf, Bangladesh	5.92	19.7	0.15	21.7	165	24.5	574

*Total TSNA*s represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).
Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosonornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNA*s = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.
Source: Stanfill et al. 2011 (4).

References

1. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf
2. Kyaing NN. Regional situation analysis of women and tobacco in South-East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://www.searo.who.int/LinkFiles/NMH_regSituationAnalysis.pdf
3. Ministry of Health and Family Welfare (Bangladesh). Global Adult Tobacco Survey: Bangladesh report 2009. Dhaka, Bangladesh: World Health Organization, Country Office for Bangladesh; 2009. Available from: http://www.who.int/tobacco/surveillance/global_adult_tobacco_survey_bangladesh_report_2009.pdf
4. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. *Tob Control*. 2011 May;20(3):e2. Epub 2010 Nov 25.

For additional information on tobacco leaf, please refer to Chapter 13: Smokeless Tobacco Use in the South-East Asia Region.

Tobacco Water

(Tuibur)

Tobacco water (also known as tuibur) is tobacco smoke-infused water that is gargled or sipped. It is commonly used in northeastern India, and is called tuibur in Mizoram and hidakpha in Manipur. This product has been used since the 19th century.^{1,2}

Common Names

Tuibur (Mizoram, India); hidakpha (Manipur, India)

Brand Names

None

Main Geographic Location

(WHO Region: Country)

South-East Asia Region: India³

Prevalence and Demographics

In 2001, tuibur was used by 7.2% of adults aged 15 and older in the Aizawl district of Mizoram, and by 6.5% of adults in the Churachandpur district of Manipur.⁴ The prevalence of use was similar among males and females.^{1,2,4}

Mode of Absorption

Oral (gargled, held in mouth)

Use Pattern

Tuibur is either sipped from a bottle or through cotton soaked with tobacco water. It is retained in the mouth or gargled for 5 to 10 minutes before it is spit out. Although people may initially use tuibur to clean their teeth, many become addicted and will use the product up to 30 times per day.^{2,4}

Main Ingredients

Tobacco smoke, water^{2,3}

Processing/Manufacturing

Cottage industry and custom-made: Tuibur is produced by passing tobacco smoke through water and is sold in glass bottles.^{1,4} Tuibur may be produced by small-scale industry or prepared at home for household use or sale.⁴

Chemical Measurements

No data available

References

1. Reddy KS, Gupta PC, editors. Report on tobacco control in India. New Delhi: Ministry of Health and Family Welfare, Government of India; 2004. Available from: http://www.who.int/ctc/reporting/Annex6_Report_on_Tobacco_Control_in_India_2004.pdf
2. World Health Organization, Regional Office of South East. Report on oral tobacco use and its implications in South East Asia. New Delhi: World Health Organization, Regional Office for South-East Asia; 2004. Available from: http://searo.who.int/en/infy/tobacco/topics/oral_tobacco_use.pdf
3. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific N-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
4. Sinha DN, Gupta PC, Pednekar M. Tobacco water: a special form of tobacco use in the Mizoram and Manipur states of India. Natl Med J India. 2004 Sep-Oct;17(5):24-57.

Tombol

Tombol is a mixture of tobacco and flavoring ingredients such as noura, slaked lime, areca nut, and catechu. It is used in Yemen and is very similar to betel quid (paan), which is used in southeast and western Asia.



Photo courtesy of Mazen Abood Bin Thabit,
University of Aden

Common Names

None

Brand Names

None

Main Geographic Location

(WHO Region: Country)

Eastern Mediterranean Region: Yemen

Prevalence and Demographics

No published information on the prevalence of tombol use is available.

Mode of Absorption

Oral (chewed, held in mouth)

Use Pattern

Tobacco and the other main ingredients are wrapped in a tombol leaf (betel leaf), placed in the mouth, and sucked and chewed. Powdered khat, a plant with psychoactive properties, may also be added.

References

Published information on tombol could not be located.

For additional information on tombol, please refer to Chapter 11: Smokeless Tobacco Use in the Eastern Mediterranean Region.

Main Ingredients

Tobacco, areca nut (fofal), noura, slaked lime (calcium hydroxide), catechu (extract from the acacia tree), tombol leaf (betel leaf), and sometimes powdered khat (*Catha edulis*) or other flavoring ingredients.

Processing/Manufacturing

Custom-made: Tombol is prepared either by a vendor or by the user, who wraps the main ingredients (tobacco, areca nut, noura, slake lime, catechu) in a tombol leaf. There are three types of tombol: (1) Sweet—a sweetening agent, usually coconut, is added to the main ingredients; (2) bitter—additives like clove oil, cardamom, and herbal medicines are used; and (3) mixed with tobacco—tombol is often mixed with either dry, thin pieces of Yemeni tobacco, called socha (similar to Indian pattiwalla), or zarda, a scented tobacco from India.

Chemical Measurements

No data available

Toombak



Photo courtesy of Clifford Watson,
Centers for Disease Control and Prevention

Toombak is a type of moist tobacco commonly used in Sudan. It is made of sun-dried ground tobacco and a solution of baking soda (sodium bicarbonate) and water.¹ The word toombak can also be used to describe the native tobacco plant used to manufacture local snuff.²

Common Names

Sauté, sute, ammari, saood

Brand Names

El-Sanf (of high quality), Wad Amari (the name of the person who introduced toombak to Sudan), Sultan El-Khaif (the master which alters the mind)

Main Geographic Locations

(WHO Region: Country)

Eastern Mediterranean Region: Sudan¹; **African Region:** Chad

Prevalence and Demographics

No recent information is available on the use of toombak. Historically, toombak was commonly used by adult Sudanese men, particularly in rural areas. The most recent published statistics (1998) found that among adults age 18 and older in the northern region of Sudan, 34% of Sudanese men and 2.5% of women use toombak.³ The prevalence of toombak use increases with age for both men and women and is more common in rural than urban areas.³

Mode of Absorption

Oral (held in mouth, sucked, used as ingredient in tombol), **Nasal**

Use Pattern

A small portion of the toombak, weighing about 10 grams, is rolled into a ball called a saffa. It is sucked slowly for 10 to 15 minutes. The saliva that is produced is then spit out by men or swallowed by women, because it is considered socially

unacceptable for women to use toombak.^{1,2} Users usually rinse their mouths with water after the saffa is removed. A regular user may use toombak as many as 10 to 20 times per day.²

Main Ingredients

Tobacco (*N. rustica*), baking soda (sodium bicarbonate, locally called atrun or natron), water^{1,2,4}

Processing/Manufacturing

Cottage industry and custom-made: Toombak tobacco leaves (*N. rustica*) are harvested and left in small heaps in a field to dry for about 45 days (sun-curing). The leaves are then tied into bundles, sprinkled with water, and stored for a few of weeks at 30 to 45° C (85–110° F) to allow fermentation. The leaves are then ground into coarse particles by toombak mills and aged for up to a year or more in burlap sacks.^{1,2} Vendors prepare ready-made toombak by gradually adding baking soda (sodium bicarbonate) to the tobacco until the mixture is approximately four parts tobacco to one part baking soda. The resulting toombak is then placed in an airtight container for about 2 hours prior to sale.^{1,2,4} While toombak is not commercially manufactured, vendors display commercial names at their shops as trademarks.

In addition to selling ready-made toombak, vendors may sell dry toombak leaves and baking soda separately so that customers can prepare their own toombak.⁴

Toombak

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNA ^s *
		mg/g wet wt					
Toombak, Sudan	7.38–10.1	9.56–28.2	5.16–10.6	14,700–516,000	115,000–368,000	4,550–6,770	295,000–992,000

*Total TSNA^s represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).
Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosonornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNA^s = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.
Source: Stanfill et al. 2011 (5).

References

1. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/index.php>
2. Idris AM, Ibrahim SO, Vasstrand EN, Johannessen AC, Lillehaug JR, Magnusson B, et al. The Swedish snus and the Sudanese toombak: are they different? *Oral Oncol.* 1998;34(6):558–66.
3. Idris AM, Ibrahim YE, Warnakulasuriya KA, Cooper DJ, Johnson NW, Nilsen R. Toombak use and cigarette smoking in the Sudan: estimates of prevalence in the Nile state. *Prev Med.* 1998;27(4):597–603.
4. Idris AM, Prokopczyk B, Hoffmann D. Toombak: a major risk factor for cancer of the oral cavity in Sudan. *Prev Med.* 1994;23:832–9.
5. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. *Tob Control.* 2011 May;20(3):e2. Epub 2010 Nov 25.

For additional information on toombak, please refer to Chapter 11: Smokeless Tobacco Use in the Eastern Mediterranean Region.

Twist

Twist, also called roll, is chewing tobacco that is twisted into rope-like strands and braided. It was popular in the United States in the late 1800s, but use of chewing tobacco began to decline with the expansion of the cigarette industry in 1918.¹ Twist is rarely used today, and sales of twist make up less than 1.0% of all smokeless tobacco sales in the United States.²



Photo courtesy of Clifford Watson,
Centers for Disease Control and Prevention

Common Names

Chew, chaw, chewing tobacco

Brand Names

Moore's Red Leaf, Cumberland, Mammoth Cave, Cotton Boll, Kentucky, Warren County, Rough Country (*American Snuff Company*)

Main Geographic Location

(WHO Region: Country)

Region of the Americas: United States¹

Prevalence and Demographics

Twist use has declined over the past century and is rare today. Although statistics specifically on the prevalence of twist use are not available, 2009 sales of twist made up 0.4% of all smokeless tobacco sales in the United States.²

Mode of Absorption

Oral (chewed, held in mouth)

Use Pattern

Users typically cut off a piece, place it in the mouth, and chew. Saliva is usually spit out, but it can also be swallowed.³

Main Ingredients

Tobacco, tobacco leaf extract, and sometimes sweetener or flavorings¹

Processing/Manufacturing

Commercial: Twist is handmade by commercial manufacturers. Air-cured or fire-cured burley tobacco leaf is treated with a tar-like tobacco leaf extract and sometimes sweeteners and other flavorings. The tobacco is then twisted into rope-like strands that are dried. The final product is a pliable, but dry, rope. Twist is sold by the piece in varying sizes, depending on the number of leaves in the twist.^{1,3}

Twist

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNAs*
		mg/g wet wt					
Twist,† United States	4.73–5.77	21.6–40.1	0.02–0.22	309–556	828–2,460	n.d.‡–104	2,590–4,950

*Total TSNAs represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).

†Twist products were commercially manufactured.

‡n.d. = not detectable.

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAs = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Lawler et al. 2013 (4).

References

1. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
2. Maxwell J. The Maxwell report: the smokeless tobacco industry in 2009. Richmond, VA: John C. Maxwell, Jr.; 2010.
3. Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR). Health effects of smokeless tobacco products. Brussels: European Commission; 2008 [cited 2012 Sept 7]. Available from: http://ec.europa.eu/health/archive/ph_risk/committees/04_scenihhr/docs/scenihhr_o_013.pdf
4. Lawler TS, Stanfill SB, Zhang L, Ashley DL, Watson CH. Chemical characterization of domestic oral tobacco products: total nicotine, pH, unprotonated nicotine and tobacco-specific *N*-nitrosamines. *Food Chem Toxicol.* 2013;57:380-6. Epub 2013 Mar 19. doi 10.1016/j.fct.2013.03.011

For additional information on twist, please refer to Chapter 9: Smokeless Tobacco Use in the Region of the Americas.

Zarda

Zarda is a flaky mixture of tobacco, lime, spices, and vegetable dyes. It is frequently chewed with chopped areca nut or used as an ingredient in paan. Zarda is commonly used in South-East Asia and in countries to which people from this region have emigrated.¹



Photo courtesy of Clifford Watson, Centers for Disease Control and Prevention

Common Names

Dokta (West Bengal, India)

Brand Names

Baba, Baghban Zafrani Zarda, Ratna Zafrani Patti, Gopal (*India*); Zahoor Zafrani Patti, Raja Jani Zafrani Patti, Sunbrand Zafrani Banarasi Patti, Shahzadi Zafrani Patti, Najma Zafran Patti (*Pakistan*); Dulal Mishti, Hakim Puri, Bat One Baba, Bullet, Surma (*Bangladesh*)

Main Geographic Locations (WHO Region: Country)

South-East Asia Region: India, Bangladesh, Myanmar, Nepal, Bhutan¹; **Eastern Mediterranean Region:** Yemen¹

Prevalence and Demographics

In India, Bangladesh, Myanmar, and Nepal, zarda is frequently used as an ingredient in paan (betel quid), particularly among middle to upper socioeconomic groups.² Although specific prevalence rates for zarda use are not available, the prevalence of smokeless tobacco use is high in these South-East Asian countries, and zarda use is common.^{1,2} Between 2007 and 2010, the adult prevalence rate of current smokeless tobacco use ranged from 18.6% to 29.6% in India, Bangladesh,³ Myanmar, Nepal, and Bhutan.⁴

Mode of Absorption

Oral (chewed, chewed in paan or tombol)

Use Patterns

Zarda may be chewed by itself, but it is usually chewed with chopped areca nuts and spices. In South-East Asia it is often used in paan, and in Yemen it is used in tombol.^{1,2,5,6}

Main Ingredients

Tobacco, lime, spices, vegetable dyes, and sometimes areca nut and/or silver flecks^{1,5}

Processing/Manufacturing

Commercial: Zarda is commercially manufactured but is usually used in user- or vendor-made paan.^{5,6} Zarda is processed by boiling broken up tobacco leaves with lime and spices until the water evaporates. It is then dried and colored with vegetable dyes.^{1,2} Zarda is sold in small packets or tins.²

Zarda

Chemical Measurements

These data are for select products and may not represent all products of this type. These data are expressed on a per gram basis for products analyzed as received. The amount absorbed will depend on the amount of product used.

Product Type*	pH	Total Nicotine	Free Nicotine	NNK	NNN	NNAL	Total TSNAst
		mg/g wet wt		ng/g wet wt			
Zarda (wet), ‡ Bangladesh	6.51	21.3	0.63	3,840	28,600	3,460	53,700
Zarda (dry), Bangladesh	6.28	9.55	0.17	457	4,280	248	9,120
Zarda, India	5.22	30.43	0.05	829	2,910	390	5,490

*All zarda products were commercially manufactured.

†Total TSNAst represent the sum of NNK, NNN, and NNAL (shown), and *N'*-nitrosoanatabine and *N'*-nitrosoanabasine (not shown).

‡This product contains areca nut.

Abbreviations: NNK = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone; NNN = *N'*-nitrosonornicotine; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; TSNAst = tobacco-specific nitrosamines; mg/g = milligram per gram; ng/g = nanogram per gram.

Source: Stanfill et al. 2011 (7).

References

1. International Agency for Research on Cancer. Smokeless tobacco and some tobacco-specific *N*-nitrosamines. IARC monographs on the evaluation of carcinogenic risks to humans. Vol. 89. Lyon, France: World Health Organization, International Agency for Research on Cancer; 2007. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol89/mono89.pdf>
2. World Health Organization, Regional Office for South-East Asia. Report on oral tobacco use and its implications in South-East Asia. New Delhi: World Health Organization, Regional Office South-East Asia; 2004. Available from: http://searo.who.int/entity/tobacco/topics/oral_tobacco_use.pdf
3. Centers for Disease Control and Prevention. Global Adult Tobacco Survey (GATS), 2008–2010: percentage of adults who currently use smokeless tobacco. Global Tobacco Surveillance System data [Internet database]. Atlanta: Centers for Disease Control and Prevention; [no date]. Available from: <http://nccd.cdc.gov/GTSSData/default/default.aspx>
4. World Health Organization. WHO report on the global tobacco epidemic: warning about the danger of tobacco. Appendix VIII, Table 8.2: crude smokeless tobacco prevalence in WHO member states. Geneva, Switzerland: World Health Organization; 2011. Available from: http://www.who.int/tobacco/global_report/2011/en_ff_global_report_2011_appendix_VIII_table_2.pdf
5. Gupta PC, Ray CS. Smokeless tobacco and health in India and South Asia. *Respirology*. 2003;8(4):419–31.
6. Reddy KS, Gupta PC, editors. Report on tobacco control in India. New Delhi: Ministry of Health and Family Welfare, Government of India; 2004. Available from: http://www.who.int/ctc/reporting/Annex6_Report_on_Tobacco_Control_in_India_2004.pdf
7. Stanfill SB, Connolly GN, Zhang L, Jia LT, Henningfield JE, Richter P, et al. Global surveillance of oral tobacco products: total nicotine, unionised nicotine and tobacco-specific *N*-nitrosamines. *Tob Control*; 2011 May;20(3):e2. Epub 2010 Nov 25.

For additional information on zarda, please refer to the following chapters: Chapter 10: Smokeless Tobacco Use in the European Region, and Chapter 13: Smokeless Tobacco Use in the South-East Asia Region.